

Volume 24

2021

**THE LIBYAN
JOURNAL
OF SCIENCE**
(An International Journal)



Published by

FACULTY OF SCIENCE
UNIVERSITY OF TRIPOLI

THE LIBYAN JOURNAL OF SCIENCE

(An International Journal)



Published by the
FACULTY OF SCIENCE
UNIVERSITY OF TRIPOLI

MEMBERS OF THE EDITORIAL BOARD

KHALED M. OUN (Editor-in-Chief)

Department of Geology, Faculty of Science,
University of Tripoli, Libya.

MOHAMED E. KELABI (Electronic Copy Coordinator)

Department of Physics, Faculty of Science,
University of Tripoli, Libya.

TAWFIK A. BULATI

Department of Mathematics, Faculty of Science,
University of Tripoli, Libya.

ALI A. EL-MAKHROUF

Department of Geology, Faculty of Science,
University of Tripoli, Libya.

RAMADAN M. ELMEHDAWI

Department of Chemistry, Faculty of Science,
University of Tripoli, Libya.

HASSAN A. MAGHRABI

Department of Zoology, Faculty of Science,
University of Tripoli, Libya.

THE LIBYAN JOURNAL OF SCIENCE

(An International Journal)

Peer reviewed international journal of science, published biannually by the Faculty of Science, University of Tripoli.

Volume 24, 2021



Published by the
FACULTY OF SCIENCE
UNIVERSITY OF TRIPOLI

Papers should be submitted to the, Editor-in-Chief, any member of the Editorial Board or the Journal Secretariat.

All correspondence should be addressed to:

Editor-in-Chief
The Libyan Journal of Science,
Faculty of Science,
University of Tripoli,
P.O.Box 13178, Tripoli-Libya.

ISSN 0368-7481 (Print)

ISSN 2521-3822 (Online)

E-mail: libyanjournal@gmail.com

Tel: 00218-21-7138882

URL: <http://libyanjournal.atSPACE.CO.UK>

Copyright © Faculty of Science, University of Tripoli, Libya

No part of this journal may be reproduced in any form without written permission from the copyright holders.

المجلة الليبية للعلوم

The Libyan Journal of Science

مجلة علمية دولية محكمة تصدر مرتين في السنة عن كلية العلوم بجامعة
طرابلس.



هيئة التحرير

أ.د. خالد محمد عون (رئيس التحرير)

أ.د. محمد إمام علي الكلابي (منسق النسخة الالكترونية)

أ.د. توفيق عبدالسلام البولاقي

أ.د. علي أحمد المخروف

أ.د. رمضان محمد المهدي

أ.د. حسن أحمد المغربي

THE LIBYAN JOURNAL OF SCIENCE

(An International Journal)

Volume 24

2021

CONTENTS

BIOLOGY

- Taher Y. Abourghiba, Mohamed T. Almskat, Alya H. Duzan
and Maha A. Elsabri, Ebtisam A. Abudaya
Study of Dermatophytosis Prevalence in Patients Attending Dermatology Clinic at
Tripoli Central Hospital..... 1
- يونس حاتم مصطفى, علي أبوالقاسم العاقل, صالح الشريف وعيسى صالح فرج
تأثير الإصابة بمرض عفن الجذور الجاف المتسبب عن فطر *Fusarium solani* على انخفاض محتوى
الكربوهيدرات بجذور الحمضيات في شمال غرب ليبيا..... 9
- Taher Y. Abourghiba, Mohammed A. Almsalati, and Zynab A. Alghadem
Investigations of the Prevalence of *Candida* Infections in Diabetes Mellitus
Patients in Tripoli..... 19
- Saleh Isslugi and Ali A. Elagael
The Effect of Mechanical Scarification on Seed Germination
of *Helianthemum lippii* (L.) Dum.Cours..... 31
- Entisar Mohamed Eljaziri and HishamAddin Bin Omar
Biology and Growth Performance for Fry Red Tilapia Feed With Spirulina..... 39
- نوال عبد السلام محفوظ و بثينة مصطفى الصغير
السُمية النباتية للزيوت العطرية الأساسية ومسحوق الأوراق لأربعة نباتات طبية على البالغات وبراغات خنفساء
Tribolium confusum Duval (Coleoptera: Tenebrionidae) المتشابهة..... 55

CHEMISTRY

- Anour Shebani, Abdalah klash, Rabia Elhabishi, Shalh Abdsalam, Wael Elhrari
Effects of Blend Composition on the UV Stability of Polyethylene Blends..... 65

MATH

- Zohra Farnana
Sobolev Spaces in Metric Spaces..... 79

PHYSICS

نجاهة سالم الشفتري سعاد محمد بوقرين، عياد الهادي الزوام، دلنده محمد ناصف و الصادق محمد القاضي دراسة خصائص نظائر $Er^{156-162}$ الشفعية-شفعية باستخدام نموذج اليوزونات المتفاعلة IBM-2.....	89
Sedig S. Farhat Use of Finite Difference Time Domain (FDTD) Technique to Calculate Poynting Vector in Free Space with Obstacles in Computational Domain.....	101
SHORT NOTE	
Entisar Mohamed Eljaziri and HishamAddin Bin Omar Effect of Feeding Fry Red Tilapia with Spirulina on Fish Health.....	119
Suggestions to Authors.....	125
تعليمات النشر.....	129

Study of Dermatophytosis Prevalence in Patients Attending Dermatology Clinic at Tripoli Central Hospital

Taher Y. Abourghiba¹, Mohamed T. Almskat, Alya H. Duzan², Maha A. Elsabri³, and Ebtisam A. Abudaya⁴

Department of Botany, Faculty of Science, University of Tripoli, Libya
¹abourghiba@hotmail.com; ²Alyaduzan@yahoo, ³Sabree1414@gamial.com, ⁴ebtisamali@yahoo.co.uk

Abstract

The current study was performed to determine the prevalence of superficial fungal infections among patients attending Dermatology Clinic at the Tripoli Central hospital. A total of 122 cases were selected with an age range 7-75 years. They were tested for fungal skin infections using microscopic and laboratory methods. The Results demonstrated that all patients had dermatophytes infections. *Tinea unguium* was the most common type of clinical infections (39.34%), followed by *Tinea capitis* (31.14%), *Tinea pedis* (22.13 %) and *Tinea faciei* (7.37 %). The results indicate that *Trichophyton violaceum* was the most common etiological agent of dermatophyte infection (59.01%) followed by *Microsporum canis* (20.49%) *Trichophyton rubrum* (12.29%), *Epidermophyton floccosum* (4.09%) and *Trichophyton menagrophitis* (3.27 %). The findings of the current study showed that higher prevalence of dermatophytes was higher in rural areas than urban areas.

Keywords: dermatophytes, Tinea infection; *Microsporum*; *Trichophyton*; *Epidermophyton*; Libya.

المستخلص

اجريت الدراسة الحالية لتحديد مدى انتشار الاصابات الفطرية السطحية بين المرضى الذين كانوا يترددون على عيادة الأمراض الجلدية بمستشفى طرابلس المركزي. تم اختيار 122 حالة تراوحت اعمارها بين (7-75 سنة) و اختبرت هذه الحالات المصابة باستخدام الفحص المجهرى للعينات والطرق المعملية. أظهرت النتائج أن جميع الحالات تعاني من اصابات فطرية جلدية وكانت سعفة الاظافر أكثر أنواع الاصابات شيوعاً (39.34%) ، تليها سعفة الرأس (31.14%) ، سعفة القدم (22.13%) وسعفة الوجه (7.37%). وظهرت نتائج هذه الدراسة اوضحت ان فطر *Trichophyton violaceum* كان من اكثر الفطريات المسببة للإصابات الجلدية بنسبة (59.01%) يليه فطر *Microsporum canis* بنسبة (20.49%) ثم فطر *Trichophyton rubrum* بنسبة (12.29%) يليه فطر *Epidermophyton floccosum* بنسبة (4.09%) وفطر *Trichophyton menagrophitis* بنسبة (3.27%) كما بينت نتائج الدراسة ان نسبة الاصابة بالأمراض الجلدية كانت اعلى في المناطق الريفية مقارنة بالمناطق الحضرية.

Introduction

Infections caused by many fungi can be divided into three groups: systemic mycosis, superficial mycosis and subcutaneous mycosis. Among superficial infection, dermatophytes is the common infection caused by fungi known as dermatophytosis. The dermatophytes are a group of fungi that are able to damage and utilize keratin found in the skin, hair, nails, horns and feathers (Grumbt et al., 2013). Evidences indicate that dermatophyte fungi are one of the most efficient human parasites, due to their efficiency in invading keratinous tissues (Zarrin et al., 2011). Dermatophytes are classified into three genera of pathogenic fungi; *Epidermophyton*, *Microsporum* and *Trichophyton*. The dermatophytosis transfers to humans from animals (zoophilic) and from soil (geophilic) or through direct infection by personal contact (Anthropophilic) (Aho, 1988). Dermatophytes have been recorded all over the world and become a public health problem in many parts of the world especially in developing countries (Nweze and Eke, 2016). Havlickova et al. (2008) and Blanka et al. (2008) have reported that 20–25% of the world's population has skin mycoses and appeared to be a frequent form of infection. Infection of nails, toes and skin by non-dermatophytes *Candida* and some molds have been increasing (Satpathi et al., 2013). The Aim of this study is to isolate and identify the fungal agents from clinical samples causing dermatophytosis in patients.

Materials and Methods

This study was conducted at the Department of Botany, Faculty of Science, University of Tripoli. The study involved patients that are clinically diagnosed for superficial mycosis. Patients data were recorded using standard format. A total of 122 samples were collected from patients attending the dermatology clinic in Tripoli Central Hospital from October 2018 to September 2019 with age range of 7-75 years. Samples were taken from skin, nails and hair by a sterile razor blade and sterile scissors. All samples were placed in labelled sterile vials and transported to the Mycology Lab. Specimens were examined and treated with 10% KOH. Specimens of nails were examined after mounting with 20% KOH for 24 hours and then examined. Each specimen was placed on a drop of KOH solution using sterile forceps and cover slip was placed on the slide.

Isolation and Culture Method

Each scraping was cultured using Sabouraud Dextrose Agar (SDA) containing 0.05 g/ml of cyclohexamide and Chloramphenicol. The plates were incubated at 25°C and were examined twice a week for perpetually up to 3 weeks. Fungal isolates were sub-cultured on the plates of SDA and were examined microscopically for morphological characteristics.

Results

In this study, the 122 scrapings collected from suspected cases of dermatophytosis, 56 cases (45.90 %) were female and 66 (54.09%) from male patients' skin infected patients.

Study of Dermatophytosis Prevalence in Patients Attending Dermatology Clinic

The etiological agents of dermatophycoses recorded in this study are *Trichophyton violacenm* was the predominant fungal species (59.01 %) followed by *Microsprum canis* (20.49 %), *Trichophyton rubrum* (12.29 %), *Epidermophyton floccosum* (4.09 %) and *Trichophyton menagrophitis* (3.27 %) (Table 1 and Fig. 1).

Table 1. Number and percentage of dermatophytes isolates

No.	Isolated species	No of isolates	In percentage
1	<i>Microsprum canis</i>	26	20.49 %
2	<i>Trichophyton menagrophitis</i>	4	3.27 %
3	<i>Trichophyton rubrum</i>	15	12.29%
4	<i>Trichophyton violacenm</i>	72	59.01%
5	<i>Epidermophyton floccosum</i>	5	4.09 %
	Total	122	100%




Fig. 1. Percentage of dermatophytes isolates

The prevalence rates of fungal infections with veneration to Tinea types Table 2 and Fig. 2) involved: Tinea capitis (31.14%), Tinea pedis (22.13 %), Tinea unguium (39.34%), and Tinea faciei (7.37%).

Table 2: Number and Percentage of Tinea types

Tinea types	Number	Percentage(%)
T. capitis	38	31.14%
T. pedis	27	22.13%
T. unguium	48	39.34%
Tinea faciei	9	7.37%




Fig. 2. Number of Tinea types

As shown in (Table 3 and Fig.3) the infection by dermatophytes was significantly higher with ages between 15- 30 years (43.4%). In contrast the percentage of infection was lower in other age groups as the following less than 15 years (12.2 %) 31-45 years(18.8 %), 46-61 years (13.9 %) and more than 61 years (11.4 %) .

Table 3. Distribution of dermatophytes fungal infections according to the age of patients.

Age groups	Frequency	Percentage
<15	15	12.2 %
15- 30	53	43.4%
31-45	23	18.8 %
46-61	17	13.9 %
>61	14	11.4 %

Results of this study have shown that the highest percentage of dermatophytes infection in the rural areas were Tinea unguium (55.73 %), Tinea capitis (27.94 %), Tinea pedis (22.05%) and Tinea faciei (8.82%) while in urban areas the fungal infections were found to be as the following Tinea unguium (37.03 %), Tinea capitis (35.18 %), Tinea pedis (22.22) and Tinea faciei (5.55 %).The number of infection was higher in rural areas compared to urban areas table 4 and 5) .

Table 4. Distribution of dermatophytes infections based on residence area (rural and urban areas)

Residence areas	Frequency	Percentage (%)
Rural areas	68	55.73 %
Urban areas	54	44.26 %
Total	122	100 %

Study of Dermatophytosis Prevalence in Patients Attending Dermatology Clinic




Fig.3. Distributions of dermatophytes fungal infections according to the ages of patients

Table 5. Distribution of the Tinea infection according patient's residence.

No	Tinea types	Urban areas		Rural areas	
		No. of patients	% of infection	No. of patients	% of infections
1	Tinea unguium	20	37.03	28	41.17
2	Tinea pedis	12	22.22	15	22.05
3	Tinea faciei	3	5.55	6	8.82
4	Tinea capitis	19	35.18	19	27.94
	Total	54	Total	68	

Discussion

In this study 122 patients suffering from dermatophytes infection were selected, in which Tinea unguium was the predominant type of infection (41.17%), Tinea capitis was the second common infection (27.94%), followed by Tinea pedis (22.05%) and Tinea faciei (8.82%). The higher prevalence of infection between toes could be attributed to the fact that they are exposed to sweating more than other parts of the body which helps the spread of the infection. The occurrence of Tinea infection was higher in 15 to 30 years (43.4%) age group followed by 31-45 years (18.8%), 46-61 years (13.9%), less than 15 years (12.2%) and >61 years (11.4%). Similar observations were reported by Sarma and Borthakur (2007) and Kansra et al. (2016). In our study, four different types of Tinea were observed among which tinea unguium was the dominant infection accounting for 39.34% of the total infections. Results of this study revealed that dermatophytes affect all ages and sexes. Similar findings were observed in previous studies (Evans and Gentles, 1985; Adefemi et al., 2011). In the current study, *Trichophyton violaceum* represents a significant percentage of dermatophyte isolates. Our finding was similar to several studies

(e.g. Woldeamanuel et al.2005; Ali-Shtayeh et al., 2002; Ameen,2010). In this study the distribution of infection by residence was 55.73 % rural areas and 44.26 % inurban areas. Although there was no significant association between infection and residence, more people in rural areas seemed to be infected than those in urban areas which may be due to prolonged exposure to soil in rural areas and have frequent encounter with animals and less hygiene standard of living. Cohen and Powdery (2004) reported that fungal infections are often associated with animals. In the present study persons of all age groups were susceptible to dermatophytosis but it appeared to be less common in age group over 61 years (11.4 % of all cases).

Conclusion and Recommendation

The present study has indicated that *Tinea unguium* was the dominant clinical infection involving 39.34% of the total cases of dermatophytosis. of the total number of patients (122). Dermatophyte isolates *Trichphyton violacem* was the predominant fungal species (59.01 %).

There is a need for further hospital survey involving large number of patients to ascertain any association between residence, age, gender and infection.

References

- Adefemi, S. A., Odeigah. L. O. and Alabi, K. M. (2011). Prevalence of dermatophytosis among primary school children in Oke-oyi community of Kwara state, Nigerian Journal of Clinical Practice, **14**(1), 23–28.
- Aho, R. (1988). Mycological studies on zoophilic dermatophyte isolates of Finish and Swedish origin Mycoses, **31**, 295–302.
- Ali-Shtayeh M. S., Salameh A. A., AbuGhdeib S. I., Jamous R. M. and Khraim, H. (2002). Prevalence of *Tinea capitis* as well as of asymptomatic carriers in school children in Nablus area (Palestine). Mycoses, **45**, 188-194.
- Ameen M. (2010). Epidemiology of superficial fungal infections. Clinics in Dermatology, **28**(2), 197–201.
- Blanka, H., Czaika V. A. and Friedrich A. (2008). Epidemiological trends in skin mycoses worldwide Mycoses, **51**, 2–15.
- Cohen, J. and Powdery, W. G. I. (2004). Infectious diseases. Mosby Edinburgh London NewYork (1): 173—179, (2) :2341-2353.
- Evans, G. V. and Gentles, J. C. (1985). Essentials of Medical Mycology. Churchill Livingstone.
- Grumbt, M., Monod, M., Yamada T., Hertweck C., Kunert, J. and Staib, P. (2013). Keratin degradation by dermatophytes relies on cysteine dioxygenase and a sulfite efflux pump, J. Invest. Dermatol., **133**, 1550–1555.
- Havlickova B., Czaika V. A. and Friedrich M. (2008). Epidemiological trends in skin mycoses worldwide. Mycoses, **51**(4), 2–15.

Study of Dermatophytosis Prevalence in Patients Attending Dermatology Clinic

- Nweze, E. I. and I. Eke, I. (2016). Dermatophytosis in northern Africa, *Mycoses*, **59**(3), 137–144.
- Satpathi, A. Achar, D. Banerjee, A. Maiti, M. Sengupta, and A. and Mohata (2013). Onychomycosis in Eastern India - study in a peripheral tertiary care centre, *Journal of Pakistan Association of Dermatologists*, 23(1),14–19, 2013
- Sarma, S. and Borthakur, A. K. (2007). A clinico-epidemiological study of dermatophytoses in Northeast India. *Indian J Dermatol Venereol Leprol.*, **73**(6), 427-8 .
- Zarrin, M. , Poosashkan, M., Mahmoudabadi , A. and Mapar, M. (2011). Prevalence of Superficial Fungal Infection in Primary School Children in Ahvaz, Iran. *Macedonian Journal of Medical Sciences*. **4**, 89-92
- Woldeamanuel Y., Leekassa R., Chryssanthou E., Menghistu Y. and Petrini B. (2005). Prevalence of tinea capitis in Ethiopian schoolchildren. *Mycoses*, **48**(2), 137–141.

تعليمات النشر

تنشر المجلة الليبية للعلوم مرتين سنويا عن طريق كلية العلوم بجامعة طرابلس وهي تهدف لنشر البحوث العلمية الاصلية في جميع افاق العلوم البحتة والتطبيقية وتقبل المجلة مخطوطات البحوث التي لم يسبق نشرها والتي ليست تحت التقييم للنشر في مكان آخر. تتم عملية المراجعة العلمية (التحكيم) عن طريق متخصصين اثنين وتحال الي مراجع ثالث في حال اختلافهما. إن من مسؤولية الكاتب أن تكون الورقة مكتوبه بلغه عربيه سليمة وتشمل ملخصا باللغة الإنجليزية (وبالإمكان الاستعانة بلجنة التحرير في اعداده).

نص المخطوط: تتم طباعة المخطوط بخط Simplified Arabic علي ورق A4 (21x29.7سم) بحيث تكون الهوامش 3 سم من اليمين واليسار ومن الأعلى والأسفل 3.8 سم لتتنسق مع شكل المجلة.
محتويات المخطوط: يطلب من الباحثين اعداد مخطوطاتهم حسب النسق التالي أو قريب منه: (1) أسم البحث، (2) أسم الباحث او البحوث كاملا وعناوينهم البريدية والإلكترونية، (3) المستخلص (عربي وانجليزي) مع كلمات دالة، (4) المقدمة، (5) الطرق والتقنيات والتجارب المستعملة او المدروسة والنطاق الجغرافي ... الخ، (6) الحسابات والتحليل والنتائج، (7) المناقشة، (8) الاستنتاجات، (9) شكر وتقدير (اختياري)، (10) المراجع و(11) الملاحق (ان وجدت).

عنوان البحث: يجب ان يكون عنوان البحث مختصرا , ومركزا في المنتصف ومكتوب بخط Simplified Arabic بحجم 16 داكن وأن يكون تحت أعلى الصفحة بسنتمترين (2سم).
الأسماء والعناوين: تدرج الأسماء الكاملة أسفل العنوان بخط Simplified Arabic حجم 11 داكن في المنتصف.

المستخلص: يجب أن يكون المستخلص متكاملًا ودالًا علي محتوى الورقة و لايزيد عن 300 كلمه، مكتوبه كفقره واحده بخط Simplified Arabic وحجم 11 داكن ويكتب العنوان "المستخلص" بخط حجم 11 داكن في منتصف السطر .

الكلمات الدالة: تكتب بنفس خط المستخلص ولا تزيد عن خمس كلمات وتوضع علي مسافة 0.5 سم عقب المستخلص مباشرة.

النص: يقسم النص الي أجزاء و اجزاء فرعيه (اذا لزم الامر) بعناوين واضحه. يبدأ النص بمقدمه (تحتوي علي المشكله والدراسات السابقة وهدف البحث) وينتهي بالاستنتاجات التي تلخص النتائج والاهداف المتحصل عليها وتوضع عناوين الاجزاء في المركز والعناوين الفرعية من اليمين وتكون جميعها بخط 14 داكن علي ان يكون النص محددًا من الجهتين .

الملاحق: توضع المعلومات الفنية التفصيلية التي قد تعوق انسياب النص في ملاحق عقب قائمة المراجع.
المراجع: إن من مسؤولية الباحث التأكد من صحة المراجع ودقتها. توضع قائمة بالمراجع التي تمت الإشارة اليها في النص وفي حال وجود مراجع لم يشر اليها سيقوم المحررون بشطبها من القائمة وتتم العودة للأصل الموجود

في التعليمات باللغة الإنجليزية Suggestions to Authors بخصوص المراجع باللغة الإنجليزية وتتم الإشارة للمراجع باللغة العربية بنفس النسق.
ملاحظة: يتم الرجوع للتعليمات باللغة الانجليزية فيما لم يتم ذكره هنا.

عنوان المجلة:

The Libyan Journal of Science
University of Tripoli
Faculty of Science
Editorial Office
P.O. Box 13503
Tripoli, Libya
E-mail: libyanjournal@gmail.com

الموقع الالكتروني للمجلة:

url: <http://libyanjournal.atspace.co.uk>

Volume 24

2021

المجلة الليبية للعلوم (مجلة دولية)



منشورات

كلية العلوم
جامعة طرابلس