

Overview Stats Comments Citations References (4) **Share**

Conference Paper **Full-text available**

### Abstract: Evaluation of vertebral fracture assessment images for the detection of abdominal aortic calcification

June 2017  
 Conference: All Wales Medical Physics and Clinical Engineering Summer Meeting 16 June 2017 - At: Village Hotel, Swansea, Wales, UK

Karima Elmasri · Yulia Hicks · Xin Yang · [Show all 6 authors](#) · William David Evans

**View your saved research**

You can access your Saved List at any time in your profile's new Saved List tab.

[OK](#)

[See details](#)

**Abstract**

Dual-energy x-ray absorptiometry (DXA) is an established modality for the assessment of bone mineral density. DXA has also been used for the detection of abdominal aortic calcification (AAC) using lateral images taken for vertebral fracture assessment (VFA). In this phantom study, the capability of VFA for the detection of AAC was investigated. A Perspex phantom of variable width in the range 15-30 cm was used to simulate abdominal soft tissue. Aluminium strips of thickness 0.05-2.0 mm were sandwiched between two halves of the phantom to mimic aortic calcification. VFA images of the phantom were acquired in single-energy mode and analysed by placing regions of interest over the aluminium strip and an adjacent area of Perspex. For each phantom width, the minimum detectable aluminium thickness was assessed visually and related to contrast-to-noise ratio (CNR). Linearity of pixel value with aluminium thickness was tested by linear regression and correlation. Repeatability was measured with five repeated scans for selected phantom configurations. The minimum thickness of aluminium that could be visualised increased with phantom width and varied from 0.05 mm at 15 cm Perspex to 0.5 mm at 30 cm Perspex; the CNR threshold was about 0.03. At all phantom widths, the variation of pixel value with aluminium thickness was strongly linear ( $r^2 > 0.98$ ,  $p < 0.001$ ). Repeatability was good with a coefficient of variation of less than 0.04%. The results of the study suggest that under idealised imaging conditions, VFA is capable of detecting small thicknesses of calcification with good linearity and repeatability.

Springer Computer Science **Ad**

**Did you know publishing a book can...?**

Connect you with your research community | Accelerate your career | Build your reputation and cement your legacy

**Have an idea for a computer science book?**

Publishing a book can help connect you with the computer science research community and accelerate your career.

[Learn more](#)

**Karima, increase the visibility of this article**

If your co-authors confirm their authorship, more researchers will find and read this article on ResearchGate.

Rebecca Jane Pettit

[Invite co-authors](#) [Not now](#)

**Related research**

**Cost effectiveness analysis of using quantitative ultrasound as a selective pre-screen for bone densitometry**

[Article](#) [Full-text available](#) February 2005

[Download](#)

[View more](#)

Public Full-text (2) Private Full-text (1)

Elmasri Abstract.pdf Content uploaded by [William David Evans](#) Author content

Page 1

**Evaluation of vertebral fracture assessment images for the detection of abdominal aortic calcification**

Karima Elmasri, Research Student  
 Yulia Hicks, Senior Lecturer  
 Xin Yang, Lecturer  
 Xianfang Sun, Lecturer  
 Rebecca Pettit, Principal Clinical Scientist  
 William Evans, Consultant Clinical Scientist  
 Cardiff University

[ElmasriKM@cardiff.ac.uk](mailto:ElmasriKM@cardiff.ac.uk)

Dual-energy x-ray absorptiometry (DXA) is an established modality for the assessment of bone mineral density [1]. DXA has also been used for the detection of abdominal aortic calcification (AAC) using lateral images taken for vertebral fracture assessment (VFA) [2]. In this phantom study, the capability of VFA for the detection of AAC was investigated.

A Perspex phantom of variable width in the range 15-30 cm was used to simulate abdominal soft tissue. Aluminium strips of thickness 0.05-2.0 mm were sandwiched between two halves of the phantom to mimic aortic calcification [3,4]. VFA images of the phantom were acquired in single-energy mode and analysed by placing regions of interest over the aluminium strip and an adjacent area of Perspex. For each phantom width, the minimum detectable aluminium thickness was assessed visually and related to contrast-to-noise ratio (CNR). Linearity of pixel value with aluminium thickness was tested by linear regression and correlation. Repeatability was measured with five repeated scans for selected phantom configurations.

The minimum thickness of aluminium that could be visualised increased with phantom width and varied from 0.05 mm at 15 cm Perspex to 0.5 mm at 30 cm Perspex; the CNR threshold was about 0.03. At all phantom widths, the variation of pixel value with aluminium thickness was strongly linear ( $r^2 > 0.98$ ,  $p < 0.001$ ). Repeatability was good with a coefficient of variation of less than 0.04%.

The results of the study suggest that under idealised imaging conditions, VFA is capable of detecting small thicknesses of calcification with good linearity and repeatability.

**References:**

- [1] A El Maghraoui and C Roux. DXA scanning in clinical practice. *Quarterly Journal of Medicine* (2008); **101**: 605–617.
- [2] K Elmasri, Y Hicks, X Yang, X Sun, R J Pettit and W D Evans. Automatic detection and quantification of abdominal aortic calcification in dual energy x-ray absorptiometry. *Procedia Computer Science* (2016); **96**: 1011–1021.
- [3] L A Fitzpatrick, A Severson, W D Edwards and R T Ingram. Diffuse calcification in human coronary arteries: association of osteopontin with atherosclerosis. *Journal of Clinical Investigation* (1994); **94**: 1597–1604.
- [4] L M Warren, A Mackenzie, D R Dance and K C Young. Comparison of the x-ray attenuation properties of breast calcifications, aluminium, hydroxyapatite and calcium oxalate. *Physics in Medicine and Biology* (2013); **58**: N103-N113.

62 of 73

[View Full-text](#)

**Similar research**

**Cost effectiveness analysis of using quantitative ultrasound as a selective pre-screen for bone densitometry**

[Article](#) [Full-text available](#) February 2005 · Technology and health care: official jo...

Victor Man Fai Sim · Michael D Stone · Ceri J Phillips · [...] · William David Evans

It has been suggested that quantitative ultrasound (QUS) could be used as a selective population pre-screen, to maximise the cost effectiveness of referral for dual energy X-ray absorptiometry (DXA) assessment of bone mineral density (BMD). We set out to...

68 Reads · 18 Citations

[Download](#) [Recommend](#) [Follow](#) [Share](#)

**NAWA NAWA Poland Ad**

**THE ULAM NAWA PROGRAMME**

A UNIQUE CHANCE TO DO YOUR OWN RESEARCH PROJECT

**Achieve your aims with the NAWA fellowship**

Are you thinking about the next step in your research career? Submit your application under the Ulam NAWA Programme!

[Find out more](#)

**Hidden Depletion of Fat-Free Mass and Bone Mineral Density in Adults With Cystic Fibrosis**

[Article](#) [Full-text available](#) December 2003 · Chest

Alina A Ionescu · William David Evans · Rebecca Jane Pettit · [...] · Dennis J Shale

Weight loss is associated with reduced survival in patients with cystic fibrosis (CF). We hypothesized that some adult patients with a normal body mass index (BMI) have evidence of hidden fat-free mass (FFM) and bone mineral density (BMD) depletion th...

25 Reads · 67 Citations

[Download](#) [Recommend](#) [Follow](#) [Share](#)

Group 1 (Mean ± SD)	Group 2 (Mean ± SD)		L4 Vertebra	L2 Vertebra
84	100			
860	624			
27.02 (9%)	16.04 (9%)	2%	27 (33.7%)	7 (8.6%)
12.04 (9%)	27.02 (9%)			
40.07 (9%)	-			
3.02 (9%)	-	1%	47 (47%)	13 (12.6%)

**Osteoporosis associated with aromatase inhibitors is different from post-menopausal osteoporosis**

[Article](#) [Full-text available](#) December 2012

Hasan Mutlu · Tuncay Aslan · Güler Silov · [...] · Abdullah Büyükçelik

osteoporosis associated with aromatase inhibitors is different from post-menopausal osteoporosis. *Adv Lab Med Int.* 2012; 3(1): 1 -5. ABSTRACT Osteoporotic effects are one of the known side effects of aromatase inhibitors that are used in the treatment ...

71 Reads · 1 Citation

[Download](#) [Recommend](#) [Follow](#) [Share](#)

**Abstract: Bone mineral density in hypothyroid and treated hyper-thyroid females on thyroxine**

[Conference Paper](#) [Full-text available](#) November 1996 · Nuclear Medicine Comm...

Rebecca Jane Pettit · William David Evans · Declan P. Coleman · [...] · John H Lazarus

It has been established that hyperthyroidism is associated with reduced bone mineral density (BMD), but it is unclear whether thyroxine therapy is also accompanied by low BMD. The aim of this study was to compare lumbar spine and femoral neck bone...

14 Reads

[Download](#) [Recommend](#) [Follow](#) [Share](#)

**The Ratio of Anterior and Posterior Vertebral Heights Reinforces the Utility of DXA in Assessment of Vertebrae Strength**

[Article](#) [Full-text available](#) May 2014 · Calcified Tissue International

Grzegorz Taton · Eugeniusz Rokita · Mariusz Korkosz · Andrzej Wrobel

The objective of the study was to introduce a new parameter describing bone strength with greater precision than the widely used antero-posterior DXA (dual-energy X-ray absorptiometry), which measures areal bone mineral density (aBMD). The adjusted...

58 Reads · 9 Citations

[Download](#) [Recommend](#) [Follow](#) [Share](#)

[View more related research](#)