

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/363857045>

Parasites of Alien fishes *Sphyraena. flavicuda* Rupplell,1838 and *Sphyraena. chrysotaenia* Klunzinger, 1884 in the Western Coast of Libya

Conference Paper · September 2022

CITATIONS

0

4 authors, including:



Aisha Seif al-naser
University of Tripoli

1 PUBLICATION 0 CITATIONS

[SEE PROFILE](#)



Sarah Ben Zeglam
University of Tripoli

2 PUBLICATIONS 0 CITATIONS

[SEE PROFILE](#)



Esmail Shakman
University of Tripoli

32 PUBLICATIONS 321 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Fish Classification [View project](#)



Gnathia sp (Gnathiidae)in some speicious of sparidae from libya [View project](#)

A. SEIFALNASER, BENZEGLAM S, SHWEHDI M, SHAKMAN E.

Tripoli University – Libya

E-mail: shugmanism@yahoo.com

**PARASITES OF ALIEN FISHES *SPHYRAENA FLAVICUDA*
RUPPLELL, 1838 AND *SPHYRAENA CHRYSOTAENIA*
KLUNZINGER, 1884 IN WESTERN COAST OF LIBYA**

Abstract

*Fish parasites are considered as a part of marine biodiversity worldwide and it is very important to know the native and invasive species in Libyan waters to be included in the Libyan marine biodiversity database. Here we investigate fish parasites in alien barracudas, *Sphyraena flavicauda* and *S. chrysotaenia*, which entered the Mediterranean through the Suez Canal. A total of 46 and 10 individuals of *S. flavicauda* and *S. chrysotaenia*, respectively were collected along the western coast of Libya. A total of seven parasites were identified. The rates of infection in *S. flavicauda* and *S. chrysotaenia* were 46% and 32% respectively, the highest prevalence was 95.7 % for the *D. cazauxi* and the lowest prevalence was 4.35% for the *Contracaecum* type III in *S. flavicauda*., Moreover, the highest prevalence in *S. chrysotaenia* was 70% for *Halacarus* sp. and the lowest prevalence was 30% for *Diplectanum dunanchae*. *D. cazauxi* and *D. dunanchae* are monogeneans, belonging to the family Diplectanidae. These results provide a first assessment of the parasitofauna of alien barracudas in the Mediterranean Sea, filling a gap of knowledge on the biological and ecological trait of these Lessepsian fishes.*

Key-words: classification, fish parasites, Lessepsian fish and Libya

Introduction

Marine ecosystems of the Mediterranean have changed at an alarming rate over the past two centuries, due to the human-mediated arrival of new species (Rilov and Galil, 2009). Marine organisms serve as hosts for various parasites and other pathogens, which can negatively affect the host population but also the food chain, with socio-economic implications (Lessios, 1988). This study aimed to investigate parasites associated to alien barracudas and assess their origin (donor region -Red Sea vs Mediterranean).

Material and methods: A total of 56 fishes were collected by local fishers along the western coast of Libya and the study was focused on metazoan parasites. The parasites examination was carried out according to Heil (2009).

Results

The rates of infection in *S. flavicauda* and *S. chrysotaenia* were 46% and 32% respectively. A total of seven parasites was identified. Their prevalence, intensity and abundance are given in Table. 1.

Tab. 2: The prevalence, intensity and abundance of parasites in *S. flavicauda* and *S. chrysotaenia* collected along the western coast of Libya

Host	<i>S. flavicauda</i>			<i>S. chrysotaenia</i>		
	Prevalence	Intensity	Abundance	Prevalence	Intensity	Abundance
<i>Hysterothylacium aduncum</i>	6.52	1	0.06	-	-	-
<i>Contracaecum type III</i>	4.35	3	0.04	-	-	-
<i>Diplectanum cazauxi</i>	95.7	1.36	0.95	-	-	-
<i>Diplectanum dunanchae</i>	85.1	1.48	0.85	30	0.30	0.30
<i>Gnathia</i> sp.	30.4	2	0.30	-	-	-
<i>halacarus</i> sp.				70	0.70	0.70
<i>Paraclanus paravus</i>	21.2	10	0.21	-	-	-

Discussion

Fish hosts can display a different parasitofauna when comparing native and invasive populations (e.g. Mele *et al.*, 2012). In this study we provide new data on the parasitofauna of alien barracudas highlighting relatively high levels of infection. Gnathiid were recorded in all examined fishes. These parasites have probably infested the exotic barracudas after their introduction in the Mediterranean Sea. Other species of parasites are known in both the Mediterranean and the Red Sea, but their short lifespan would indicate that they were acquired in the newly invaded habitat (Mackenzie and Aباunza, 2014).

Bibliography

- HEIL N., (2009) - National Wild Fish Health Survey – Laboratory Procedure Manual, fifth ed. US Fish and Wildlife Service, Warm Springs
- LESSIOS H.A. (1988) - Mass mortality of *Diadema antillarum* in the Caribbean: what have we learned? *Ann Rev Ecol Syst* 19: 371-393.
- MACKENZIE K., ABAUNZA P. (2014) - Parasites as biological tags. In *Stock Identification Methods* (pp. 185-203). Academic Press
- MELE S., MACIAC D., GOMEZ-VIVES M. J., GARIPPA G., ALEMANY F., MERELLA P. (2012) - Metazoan parasites on the gills of the skipjack tuna *Katsuwonus pelamis* (Osteichthyes: Scombridae) from the Alboran Sea (western Mediterranean Sea). *Diseases of aquatic organisms*, 97(3), 219-225.
- RILOV G., GALIL B. (2009) - Marine bioinvasions in the Mediterranean Sea—history, distribution and ecology. In *Biological invasions in marine ecosystems* (pp. 549-575). Springer, Berlin, Heidelberg.