

Chloroform Fraction from Methanol Extract of Starfish *Acanthaster planci* Stimulates Catfish (*Clarias sp.*) Macrophage Immunomodulatory Activity

Janib Achmad¹, Ervia Yudiati^{2*}, Nuril Azhar², Rabia Alghazeer³

¹Faculty of Fisheries and Marine Science, University of Khairun Ternate
Jl. Pertamina Kampus 2, Kel. Gambesi, Ternate Selatan Indonesia

²Department of Marine Science, Faculty of Fisheries and Marine Science, Diponegoro University
Jl. Prof. Jacub Rais, Tembalang Semarang, Jawa Tengah 50275 Indonesia

³Department of Chemistry, Faculty of Sciences, University of Tripoli
Tariq Sayyidi al Misri Road, Tripoli, Libya
Email: eyudiati@gmail.com

Abstract

Catfish (*Clarias sp.*) is Indonesia's most important and popular freshwater commodity, widely cultured and intensively grown. This study is aimed to find out the effects of the chloroform fraction from a methanol extract of *Acanthaster planci* on non-specific immunity in *Clarias sp.* Thin Layer Chromatography analysis of the methanol extract's chloroform fraction yielded three fractions: FrK13.1, FrK13.2, and FrK13.3. The FrK13.3 fraction of *Acanthaster planci* had a significant influence on Phagocytic Activity at doses of 0.1 (54.09 ± 8.99 %), 0.3 (48.16 ± 3.34 %), 0.5 (50.39 ± 5.71 %), and 0.7 mg/kg Body Weight (46.58 ± 0.37 %). At 0.5 mg/kg Body Weight (4.03 ± 1.40 %), the Phagocytic Index of FrK13.2 was significantly higher than the control. At 0.5 mg/kg Body Weight (38.22 ± 1.43 cell 10⁶/ml) and 0.7 mg/kg BW (40.41 ± 1.92 cell 10⁶/ml), the number of leukocytes/white blood cells was significantly higher than the control. Nitroblue Tetrazolium and Total Plasma Protein were not significantly different from the control. Based on the results, the FrK13.3 was leading on phagocytic capacity and number of leukocytes

Keyword: *Acanthaster planci*, *Clarias sp.*, chloroform fraction, non-specific immunity, macrophage

INTRODUCTION

Catfish (*Clarias sp.*) is Indonesia's most important freshwater commodity (Gustiano *et al.*, 2021). Some cultivators are willing to increase production yield by implementing an intensive cultivation system and rearing fish in a small area with a high stocking density. *Clarias sp.* culture has some deserving perspectives in Indonesia (Diatin *et al.*, 2021; Gustiano *et al.*, 2021; Henriksson *et al.*, 2017). *Clarias sp.* is rich in nutrition, such as protein (16%), lipid (16.92%), fatty acid (24.51%) palmitic acid (29.57%), oleic acid, and 19.07% linoleic acid. Moreover, *Clarias sp.* is also have a valuable amino acids, specifically 13,818% essential amino acids (0.796% Histidine; 1.708% Threonine; 1.719% Valine; 1.205% Methionine; 2.613% Lysine; (Yu *et al.*, 2017). However, high stocking density in an intensive system will lead fish to become stressed, resulting in a decline in fish immunity. As a result, the fish will be susceptible to the disease (Pietrzak *et al.*, 2020; Purbomartono *et al.*, 2019).

Clarias sp. is susceptible to diseases caused by pathogen bacteria such as *Aeromonas hydrophyla* (Sellegounder *et al.*, 2018), *Flavobacterium columnare* (Harikrishnan *et al.*, 2018), and Channel catfish virus (Boon *et al.*, 1988). Chemical substances may effectively control pathogen, but the administration over a long-term period give notice of being harmful (Barman *et al.*, 2013; Chiu *et al.*, 2008; Chopra *et al.*, 2000; Purbomartono *et al.*, 2019; Yang *et al.*, 2018). Pathogenic bacteria can become resistant to chemical substances found in the water environment. Furthermore, the chemical substances' residues can be harmful to human health. By increasing the fish's immunity with an immunomodulator or natural chemical compounds, hazardous substances can be avoided. Furthermore, immunomodulator from natural product is environmentally friendly and does not leave any chemical residues that are harmful to human health (A. Cheng *et al.*, 2008; Jensch-Junior *et al.*, 2006; Magnadóttir, 2010; Magnadóttir, 2006).

According to Chiu *et al.* (2008), the use of improper immunomodulators preserves the stressful conditions that negatively impact fish physiological conditions. It has been widely reported that

natural substances such as polysaccharide polymers, alginates, fucoidan from seaweed, and glucoside from sea cucumbers can boost fish and shrimp immunity (A. Cheng *et al.*, 2008; W. Cheng dan Yu, 2013; Fuchs *et al.*, 2015; Purbomartono *et al.*, 2019; Yudiati *et al.*, 2019). El-Boshy *et al.* (2014) discovered that dietary fucoidan increased non-specific immune response and disease (*A. hydrophila*) resistance in *Clarias gariepinus*. Another study found that *Clarias gariepinus* could withstand a *Flavobacterium columnare* challenge when fed an *Agaricus bisporus*-enriched diet (Harikrishnan *et al.*, 2018).

The primary mechanism of the fish immune response is from aquatic organisms, which includes specific and non-specific responses such as memory, specificity, and recognition of foreign substances (Cheng and Yu, 2013; Mehana *et al.*, 2015; Pietrzak *et al.*, 2020). In terms of macrophage immunomodulatory activity, our previous study demonstrated the effectiveness of *Acanthaster planci* in the chloroform fraction of methanol extract (M. J. Achmad *et al.*, 2018). In fact, in nature, *A. planci* is a dangerous coral destroyer. Pratchett *et al.* (2009) found *A. planci* densities of 162 sea stars per hectare in Papua New Guinea, causing extensive coral mortality. This outbreak killed up to 55% of live corals, reducing overall coral cover from 42.4 % in 2005 to 19.1 % in March 2006. There must be some interesting compound in this particular Sea Star that needs to be explored. As a result, the use of *A. planci* as a potential immunostimulant must be exposed and assessed. This study is aimed to find out the immunomodulatory activities of three active fractions extracted from *A. planci*, namely FrKI3.1, FrKI3.2, and FrKI3.3, and its modulation of *Clarias* sp. non-specific immunity.

MATERIAL AND METHODS

A. planci was collected on Ternate Island, North Maluku, Indonesia. Ethyl acetate 99.5% (Sigma, USA), Chloroform 99% (Merck, USA), Methanol 95% (Merck, USA), NaOH 99% (Merck, USA), Silica gel 60 F254 plates (Merck, USA), preparative silica gel GF254 (Merck, USA), C18 (Reversed-phase, Pyrex USA), and KLT (E. Merck) plates were used as solvents and purification materials. HCl 37 % (Merck, USA), diethyl ether 99.7% (Merck, USA), acetone 99% (Merck, USA), ethanol 96% (Merck, USA), Calcium Chloride 99% (Sigma, USA), 100 mg/ml Bovine Serum Albumin/BSA 99% (Sigma, USA), test kit protein, Biorad methods (Sigma, USA), and fish blood samples were used in the haematology test. For the fish diet, commercial fish feed (Pokphand 781-2-a) was used.

Isolation and Purification of Bioactive Compound *A. planci*

Extraction and partition were carried out following the method used by M. Achmad *et al.* (2020); M. J. Achmad *et al.* (2018); M. J. Achmad *et al.* (2014). Methanol was used to extract and filter *A. planci* samples. To obtain the methanol extract, a rotative evaporator (Rotavapor® R-100, Buchi) was used at a temperature of 35°C. By separatory funnel, the methanol extract was partitioned to chloroform, hexane, and water in a 1:1:1 ratio. Partitioning was performed to obtain three fractions: hexane, chloroform, and water. After that, the fractions were evaporated. Since the water and hexane fractions were done in our previous study, in this present study we continue with the chloroform fraction (M. J. Achmad *et al.*, 2018). The chloroform fraction was then fractionated using eluent Chloroform: Methanol. This fractionation was then chemically profiled on a 5 cm column using Choloform: Methanol = 8:2 (10 ml) as eluent. FrKI3 fractions were chosen for hematological testing based on the results of the chemical profiling process.

Administration of Fraction FrKI3 *A. planci*

During fish cultivation, Pokphand 781-2-a commercial feed was administered. Catfish were raised in a fiber tank and feeding was done twice a day on an ad libitum basis. The average size of the fish was 40 cm long and 30 cm wide. The injection method was used to administer FrKI3 fraction to fish, and it was repeated three times. Three fish were injected with FrKI3 fractions at 0.1 mg/kg BW, 0.3 mg/kg BW, 0.5 mg/kg BW, and 0.7 mg/kg BW in each treatment. Three days after the first injection, the injections were repeated. The blood draw occurred after the second injection and on the fifth day following the second injection.

Hematology Test

Immunomodulatory activity of fish injected with FrK13 fractions to non-specific defenses was assessed using hematological parameters such as macrophage phagocytosis, number of leukocytes, nitroblue tetrazolium (NBT), total plasma protein, and leukocyte differentiation (monocytes, neutrophils, and lymphocytes), as well as physiological parameters (hematocrit and leucocyte). The hematological parameters were observed using the techniques followed by M. J. Achmad *et al.* (2014); A. Cheng *et al.* (2008).

Phagocytosis Test

Blood fish sample was carried out using a syringe which was previously coated with EDTA solution and the sample was then transferred into a microtube, immediately. The blood was then drawn using a capillary tube and centrifuged at 1500 rpm within 5 minutes. Afterward, the capillary tube was cut to take the white blood cells to be placed into a microtube. A total of 50 μ l of white blood was inserted into the well of microtiter plate 96. Furthermore, the antigen *Staphylococcus* sp. with a volume of 50 μ l was mixed into a microtiter plate containing white blood and mixed. Subsequently, it was incubated for 30 minutes and stained with 10% Giemsa solution in ethanol. 5 μ l of incubated white blood was taken out using a micropipette and then dropped on a glass preparation to be examined. Before this, absolute methanol was used for fixation, and then dried, and followed by staining with 10% Giemsa solution for 15 minutes. The observation of macrophages was conducted using a microscope with 1000 times magnification, and the results of the observations were analyzed to measure the phagocytic activity (PA) and phagocytic index (PI).

White Blood Cell (WBC) Test

The calculation of white blood cells (leukocytes) was done by taking \pm 50 μ l of fish blood. Blood samples were taken using 0.5 of a hemocytometer tube and added with 1N HCl. It was then mixed before being observed with a hemocytometer under a microscope to calculate the number of white blood cells.

Total Plasma Protein Test

The measurement of total plasma protein was conducted by filling the microtube with 798 μ l aqua bides according to the number of samples. Each microtube was added with 2 μ l serum and a 200 μ l protein test kit (Biorad) before being mixed using the pipette, and followed 15 minutes incubation. The absorbance was measured using a spectrophotometer at a wavelength of 595-610 nm, and the concentration of total plasma protein was then determined using a standard curve.

Nitrobluetetrazolium (NBT)

100 μ l of fish blood was put into a microtube and added with 0.2% NBT solution at a similar volume. The mix was incubated for 30 minutes at 30°C. After the incubation, a 50 μ l sample was then put into 1 ml N-N dimethylformamide solution and then centrifuged for 5 minutes at 3000 rpm. The supernatant was observed, and its absorbance was measured using a spectrophotometer at a wavelength of 595-610 nm.

Statistical Analysis

The obtained data were tested for variance using analysis of variance (ANOVA), using R Studio application. If the analysis of the variance test revealed a significant difference, it was further tested using Tukey HSD multiple comparisons of means with a 95 % family-wise confidence level ($P < 0.05$). NBT, PA, PI, total plasma protein, hematocrit, leucocrit, number of SDPs, and % of leukocyte types among the variables were examined.

RESULT AND DISCUSSION


Chemical Profiling

Table 1 shows the results of fractionation with four elutions: CHCl₃: MeOH = 2:1, CHCl₃: MeOH = 2:2, CHCl₃: MeOH = 1:2, and MeOH elution. FrK13.1; FrK13.2; and FrK13.3, as a combination of several

fractions (F1-F6, F7-F15, and F16-F25), respectively, comprised the fractionation in 25 fractions. This classification is based on the similarity of retention factors (RF), and the combination of these fractions results in chemical profiling from the TLC plate. The first fraction (FrKI3.1) was developed from six fractions namely F1, F2, F3, F4, F5, and F6. The second fraction (FrKI3.2) was developed from nine fractions, namely F7, F8, F9, F10, F11, F12, F13, F14, and F15. The last third fraction (FrKI3.3) was developed from 10 fractions F16, F17, F18, F19, F20, F21, F22, F23, F24, and F25. FrKI3.1; FrKI3.2, and FrKI3.3 were then continued with chemical profiling with static phase SiO₂ 60 F₂₅₄: mobile phase n-Hx: EtOAc = 9 :1 v/v. The spot developed was detected under UV light at 254 nm. The R_f value from major to minor of FrKI3.1; FrKI3.2 and FrKI3.3 were 0.7; 0.5; and 0.4, respectively. The major spot developed from FrKI3.1, followed by FrKI3.2., and the FrKI3.3 developed spot was minimum.

The differences in R_f value are based on the distance of the spot developed in the TLC plate. The R_f value of each compound is related to the polarity. In comparison with the polar compound, the nonpolar compound will move more distance on a plate, so, therefore leads to a higher R_f value. In contrast to the lower compound polarity, the higher compound polarity has a stronger interaction with silica gel (static phase) rather than the lower ones (Fair & Kormos, 2008; Reich & Schibli, 2007; Stoddard *et al.*, 2007). The chemical profiling in this present study is to monitor the fraction movement, identify the existing compound in the crude extract, and to define the compound purity (Reich & Schibli, 2007; Stoddard *et al.*, 2007). Hematological tests were then performed on the three fractions to get the best fraction.

Table 1. Chemical profiling of the FrKI3 fraction resulted in three (3) subfractions (FrKI3.1; FrKI3.2; and FrKI3.1)



Remarks: stationary phase of SiO₂ 60 F₂₅₄: mobile phase of n-Hx: EtOAc = 9:1 v/v, detected under UV 254 nm.

Phagocytic Activity (PA) and Phagocytic Index (PI)

Figure 1 depicts the results of the phagocytic activity in three fractions of FrKI3. FrKI3.1 (Figure 1A) significantly exceeded control at doses of 0.3 (51.11±5.47 %) and 0.5 (50.96±0.64 %) mg/kg BW, whereas FrKI3.2 exceeded control (36.3±7.47 %) at a dose of 0.7 mg/kg BW. Furthermore, the PA value of FrKI3.3 significantly exceeded the control (30.99±4.51 %) at all doses, i.e., 46.58-54.09 % with a standard deviation of 0.37-8.99 %. Only a dose of 0.5 mg/kg BW FrKI3.2 (4.03±1.40 %) significantly outperformed the control (1.69±0.32 %) in the PI value (Figure 1B).

The chloroform fraction from the methanol extract of *A. planici* stimulated the nonspecific fish immune system in terms of phagocytic capacity (Phagocytic Activity and Index) and the number of leukocytes/white blood cells (WBC). At 0.3-0.5; 0.7 mg/kg BW and all doses, these three fractions of FrKI3.1, FrKI3.2, and FrKI3.3 had a significant effect on Phagocytic Activity. PA indicates an increase in cellular response, which eventually leads to an increase in humoral response (M. J. Achmad *et al.*, 2018; M. J. Achmad *et al.*, 2014; Jensch-Junior *et al.*, 2006; Yudiati *et al.*, 2019). The PI (Phagocyte Index) value was significantly higher than the control, FrKI3.2 at a dose of 0.5 mg/kg BW (4.03±1.40).




Figure 1. Phagocytic Activity (a) and Phagocytic index (b) of FrKI3.1; FrKI3.2; and FrKI3.3 on *Clarias* sp., analyzed with two-way ANOVA (R Studio Application). The significant differences were indicated by different notations. Phagocytic activity and index were significant in the dose and not significant in fractions. Only phagocyte activity had an interaction between dose and fraction.

According to Magnadóttir (2006); Purbomartono *et al.* (2019), if the PI value in the treatment group is higher than the control group, it indicates a stimulatory effect on increasing PI due to the active compound from the material test. In line with previous research, the hexane fraction of the methanol extract had a significant effect on Phagocyte Activity and Index (M. J. Achmad *et al.*, 2018).

Number of Leukocytes/White Blood Cells (WBC)

The results of total leukocytes in three fractions of FrKI3 are presented in Figure 2. The FrKI3.1 fraction did not differ significantly from the control ($p > 0.05$). FrKI3.2 at 0.5 mg/kg BW (38.22 ± 1.43 cell. 10^6 /ml) and FrKI3.3 at 0.7 mg/kg BW (40.42 ± 1.92 cell. 10^6 /ml) were significantly different from the FrKI3.2 control (21.34 ± 2.25 cell. 10^6 /ml) and FrKI3.3 control (22.63 ± 3.00 cell. 10^6 /ml, respectively). The number of leukocytes/white blood cells (WBC) significantly exceeded the control.

Fish health, environmental conditions, and fish species all influence the number of leukocytes (M. J. Achmad *et al.*, 2018; Chiu *et al.*, 2008; Jensch-Junior *et al.*, 2006; Purbomartono *et al.*, 2019; Yudiati *et al.*, 2019). Le *et al.* (2016) discovered two new pyrrole oligoglycosides, plancipyrosides A and B (1 and 2), in methanol extract from *A. planci* Macrophages. Macrophages in RAW264.7 was able to suppress Lipopolysaccharide-Induced Nitric Oxide Production. NO is a signaling molecule that regulates physiological processes (such as vasodilation and neurotransmission) as well as the inflammatory response (Pacher *et al.*, 2007). It has cytotoxic properties against microbial pathogens but can also be harmful to host tissues (Korhonen *et al.*, 2005). Prevention of NO overproduction in living organisms via regulatory pathway control can aid in the treatment of high NO-mediated disorders without changing the physiological level of the NO radical point (Aktan, 2004).

Nitroblue Tetrazolium (NBT)

The results of NBT in three fractions of FrKI3 are presented in Figure 3. All FrKI3 fractions were not significantly different from the control ($p > 0.05$). The respiratory level of macrophage cells was tested during the phagocytosis process using nitro-blue tetrazolium (NBT) solution, which was reduced into formazan when the reaction occurred with the radical oxygen produced by macrophage (Fuchs *et al.*, 2015; Nieves-Rodríguez *et al.*, 2018; Wang *et al.*, 2017). During phagocytosis, the macrophage frequently performed a respiratory burst, resulting in reactive oxygen species (ROS), which play an important role in pathogen defense. NADPH oxidase was a key enzyme in this process (Fuchs *et al.*, 2015; Nieves-Rodríguez *et al.*, 2018; Vetvicka *et al.*, 2013; Wang *et al.*, 2017).




Figure 2. Total leukocytes of FrKI3.1; FrKI3.2; and FrKI3.3 on *Clarias* sp., analyzed with two-way ANOVA (R Studio Application). The significant differences were indicated by different notations. Leucocyte was significant in dose and fractions. Leucocytes have no interaction between dose and fraction.




Figure 3. Values of nitroblue tetrazolium (NBT) of FrKI3.1; FrKI3.2; and FrKI3.3 on *Clarias* sp. were analyzed with two-way ANOVA (R Studio Application). The significant differences were indicated by different notations.




Figure 4. TPP of FrKI3.1; FrKI3.2; and FrKI3.3 on *Clarias* sp. were analyzed with two-way ANOVA (R Studio Application). The significant differences were indicated by different notations.

The best dose of NBT on *Clarias batrachus* and rainbow trout fish was 0.2-0.4 mg/kg body weight. Fish had a killing mechanism by phagocytosis cells through the production of radical faculae lysosome that could increase the permeability of bacterial cells, allowing fluid to enter the cells and cause plasmolysis (Fuchs *et al.*, 2015; Guzmán-Villanueva *et al.*, 2014).

Total Protein Plasma (TPP)

Results of TPP in three fractions of FrKI3 are presented in Figure 4. All FrKI3 fractions were not significantly different from the control ($p>0.05$). Total protein plasma may also contribute to an increase in fish immunity to disease. Furthermore, total protein plasma may indicate the amount of nitrogen in amino acids (Fuchs *et al.*, 2015; Wang *et al.*, 2017).

CONCLUSION

The methanol extract's chloroform fraction yielded three fractions: FrKI3.1, FrKI3.2, and FrKI3.3. The best PA was gained in FrKI3.3 of *Acanthaster planci* fraction at the dose at doses of 0.1 mg/kg

BW. The PI, and number of leukocytes/white blood cells, the FrkI3.2 shows the best results. Eventhough, the Nitroblue Tetrazolium and Total Plasma Protein were not significantly different from the control. This present research shows that fractionation can stimulate Macrophage Immunomodulatory Activity even at low doses.

REFERENCES

- Achmad, M., Isnansetyo, A., Andriani, R., Samman, A., & Marus, I. (2020). The analysis of challenges test of catfish (*Clarias* sp.) with fatty acid compounds from starfish (*Acanthaster planci*). *IOP Conference Series: Earth and Environmental Science*, 584, p.012014. doi:10.1088/1755-1315/584/1/012014
- Achmad, M.J., Isnansetyo, A., & Kasanah, N. (2018). Macrophage Immunomodulatory Activity of Unsaturated Fatty Acid Isolated from the Crown-of-thorns Star Fish (*Acanthaster planci*). *Pharmacognosy Journal*, 10(5), 951-957. doi: 10.5530/pj.2018.5.161
- Achmad, M. J., Isnansetyo, A., Kasanah, N., Ustad, U., & Kamiso, K. (2014). Immunostimulatory effect of fatty acid from star fish (*acanthaster planci*) on lymphocyte proliferation in-vitro. *Squalen Bulletin of Marine Fisheries Postharvest Biotechnology and Bioprocess Engineering*, 9(3), 107-114. doi: 10.15578/squalen.v9i3.109
- Aktan, F. (2004). iNOS-mediated nitric oxide production and its regulation. *Life Sciences*, 75(6), 639-653. doi: 10.1016/j.lfs.2003.10.042
- Barman, D., Nen, P., Mandal, S., & Kumar, V. (2013). Immunostimulants for Aquaculture Health Management. *Journal of Marine Science: Research & Development*, 3(3), 1-11. doi: 10.4172/2155-9910.1000134
- Boon, J.H., McDowell, T., & Hedrick, R.P. (1988). Resistance of the African (*Clarias gariepinus*) and the Asian catfish (*Clarias batrachus*) to channel catfish virus. *Aquaculture*, 74(3), 191-194. doi: 10.1016/0044-8486(88)90362-6
- Cheng, A., Wan, F., Wang, J., Jin, Z., & Xu, X. (2008). Macrophage immunomodulatory activity of polysaccharides isolated from *Glycyrrhiza uralensis* Fish. *International Immunopharmacology*, 8(1), 43-50. doi: 10.1016/j.intimp.2007.10.006
- Cheng, W., & Yu, J.S. (2013). Effects of the dietary administration of sodium alginate on the immune responses and disease resistance of Taiwan abalone, *Haliotis diversicolor supertexta*. *Fish & Shellfish Immunology*, 34(3), 902-908. doi: 10.1016/j.fsi.2012.12.020
- Chiu, S.T., Tsai, R.T., Hsu, J.P., Liu, C.H., & Cheng, W. (2008). Dietary sodium alginate administration to enhance the non-specific immune responses, and disease resistance of the juvenile grouper *Epinephelus fuscoguttatus*. *Aquaculture*, 277(1), 66-72. doi: 10.1016/j.aquaculture.2008.01.032
- Chopra, A.K., Xu, X., Ribardo, D., Gonzalez, M., Kuhl, K., Peterson, J.W., & Houston, C.W. (2000). The cytotoxic enterotoxin of *Aeromonas hydrophila* induces proinflammatory cytokine production and activates arachidonic acid metabolism in macrophages. *Infection and Immunity*, 68(5), 2808-2818. doi: 10.1128/iai.68.5.2808-2818.2000
- Diatin, I., Shafruddin, D., Hude, N., Sholihah, M. a., & Mutsmir, I. (2021). Production performance and financial feasibility analysis of farming catfish (*Clarias gariepinus*) utilizing water exchange system, aquaponic, and biofloc technology. *Journal of the Saudi Society of Agricultural Sciences*, 20(5), 344-351. doi: 10.1016/j.jssas.2021.04.001
- El-Boshy, M., El-Ashram, A., Risha, E., Abdelhamid, F., Zahran, E., & Gab-Alla, A. (2014). Dietary fucoidan enhance the non-specific immune response and disease resistance in African catfish, *Clarias gariepinus*, immunosuppressed by cadmium chloride. *Veterinary Immunology and Immunopathology*, 162(3), 168-173. doi: 10.1016/j.vetimm.2014.10.001
- Fair, J.D., & Kormos, C.M. (2008). Flash column chromatograms estimated from thin-layer chromatography data. *Journal of Chromatography A*, 1211(1-2), 49-54. doi: 10.1016/j.chroma.2008.09.085
- Fuchs, V.I., Schmidt, J., Slater, M.J., Zentek, J., Buck, B.H., & Steinhagen, D. (2015). The effect of supplementation with polysaccharides, nucleotides, acidifiers and *Bacillus* strains in fish meal and soy bean based diets on growth performance in juvenile turbot (*Scophthalmus maximus*). *Aquaculture*, 437, 243-251. doi: 10.1016/j.aquaculture.2014.12.007

- Gustiano, R., Prakoso, V., Radona, D., Dewi, R., Saputra, A., & Nurhidayat, N. (2021). A sustainable aquaculture model in Indonesia: multi-biotechnical approach in *Clarias* farming. *IOP Conference Series: Earth and Environmental Science*, 718, p.012039. doi: 10.1088/1755-1315/718/1/012039
- Guzmán-Villanueva, L.T., Ascencio-Valle, F., Macías-Rodríguez, M.E., & Tovar-Ramírez, D. (2014). Effects of dietary β -1,3/1,6-glucan on the antioxidant and digestive enzyme activities of Pacific red snapper (*Lutjanus peru*) after exposure to lipopolysaccharides. *Fish Physiology and Biochemistry*, 40(3), 827-837. doi: 10.1007/s10695-013-9889-0
- Harikrishnan, R., Naafar, A., Musthafa, M.S., Ahamed, A., Arif, I.A., & Balasundaram, C. (2018). Effect of *Agaricus bisporus* enriched diet on growth, hematology, and immune protection in *Clarias gariepinus* against *Flavobacterium columnare*. *Fish and Shellfish Immunology*, 73, 245-251. doi: 10.1016/j.fsi.2017.12.024
- Henriksson, P.J.G., Tran, N., Mohan, C.V., Chan, C.Y., Rodriguez, U.P., Suri, S., Mateos, L.D., Utomo, N.B.P., Hall, S. & Phillips, M.J., (2017). Indonesian aquaculture futures – Evaluating environmental and socioeconomic potentials and limitations. *Journal of Cleaner Production*, 162, 1482-1490. doi: 10.1016/j.jclepro.2017.06.133
- Jensch-Junior, B.E., Pressinotti, L. N., Borges, J.C.S., & da Silva, J.R.M.C. (2006). Characterization of macrophage phagocytosis of the tropical fish *Prochilodus scrofa* (Steindachner, 1881). *Aquaculture*, 251(2), 509-515. doi: 10.1016/j.aquaculture.2005.05.042
- Korhonen, R., Lahti, A., Kankaanranta, H., & Moilanen, E. (2005). Nitric Oxide Production and Signaling in Inflammation. *Current Drug Targets - Inflammation & Allergy*, 4(4), 471-479. doi: 10.2174/1568010054526359
- Le, T.V., Hanh, T.T.H., Huong, P.T.T., Dang, N.H., Van Thanh, N., Lyakhova, E., Cuong, N.X., Nam, N.H., Van Kiem, P., Kicha, A. and Van Minh, C., 2016. Pyrrole oligoglycosides from the starfish *Acanthaster planci* suppress lipopolysaccharide-induced nitric oxide production in RAW264. 7 macrophages. *Chemical and Pharmaceutical Bulletin*, 64(11), 1654-1657. doi: 10.1248/cpb.c16-00585
- Magnadottir, B. (2010). Immunological control of fish diseases. *Marine biotechnology*, 12(4), 361-379. doi: 10.1007/s10126-010-9279-x
- Magnadóttir, B. (2006). Innate immunity of fish (overview). *Fish and Shellfish Immunology*, 20(2), 137-151. doi: 10.1016/j.fsi.2004.09.006
- Mehana, E.S., Rahmani, A., & Aly, S. (2015). Immunostimulants and Fish Culture: An Overview. *Annual Research and Review in Biology*, 5, 477-489. doi: 10.9734/ARRB/2015/9558
- Nieves-Rodríguez, K.N., Álvarez-González, C.A., Peña-Marín, E.S., Vega-Villasante, F., Martínez-García, R., Camarillo-Coop, S., Tovar-Ramírez, D., Guzmán-Villanueva, L.T., Andree, K.B. & Gisbert, E. (2018). Effect of β -Glucans in Diets on Growth, Survival, Digestive Enzyme Activity, and Immune System and Intestinal Barrier Gene Expression for Tropical Gar (*Atractosteus tropicus*) Juveniles. *Fishes*, 3(3), p.27. doi: 10.3390/fishes3030027
- Pacher, P., Beckman, J.S., & Liaudet, L. (2007). Nitric Oxide and Peroxynitrite in Health and Disease. *Physiological Reviews*, 87(1), 315-424. doi: 10.1152/physrev.00029.2006
- Pietrzak, E., Mazurkiewicz, J., & Slawinska, A. (2020). Innate Immune Responses of Skin Mucosa in Common Carp (*Cyprinus Carpio*) Fed a Diet Supplemented with Galactooligosaccharides. *Animals*, 10(3), p.438. doi: 10.3390/ani10030438
- Pratchett, M.S., Schenk, T.J., Baine, M., Syms, C., & Baird, A.H. (2009). Selective coral mortality associated with outbreaks of *Acanthaster planci* L. in Bootless Bay, Papua New Guinea. *Marine Environmental Research*, 67(4-5), 230-236. doi: 10.1016/j.marenvres.2009.03.001
- Purbomartono, C., Isnansetyo, A., Murwantoko, & Triyanto. (2019). Dietary Fucoidan from *Padina boergesenii* to Enhance Non-specific Immune of Catfish (*Clarias* sp.). *Journal of Biological Sciences*, doi: 10.3923/jbs.201
- Reich, E., & Schibli, A. (2007). High-performance thin-layer chromatography for the analysis of medicinal plants. Thieme. FAO
- Sellegounder, D., Gupta, Y.R., Muruganankumar, R., & Senthilkumaran, B. (2018). Enterotoxic effects of *Aeromonas hydrophila* infection in the catfish, *Clarias gariepinus*: Biochemical, histological

- and proteome analyses. *Veterinary Immunology and Immunopathology*, 204, 1-10. doi: 10.1016/j.vetimm.2018.08.008
- Stoddard, J.M., Nguyen, L., Mata-Chavez, H., & Nguyen, K. (2007). TLC plates as a convenient platform for solvent-free reactions. *Chemical Communications*, (12), 1240-1241. doi: 10.1039/b616311d
- Vetvicka, V., Vannucci, L., & Sima, P. (2013). The Effects of β - Glucan on Fish Immunity. *North American Journal of Medical Sciences*, 5(10), 580-588. doi: 10.4103/1947-2714.120792
- Wang, W., Sun, J., Liu, C., & Xue, Z. (2017). Application of immunostimulants in aquaculture: current knowledge and future perspectives. *Aquaculture Research*, 48(1), 1-23. doi: 10.1111/are.13161
- Yang, P., Hu, H., Liu, Y., Li, Y., Ai, Q., Xu, W., Zhang, W., Zhang, Y., Zhang, Y. & Mai, K., (2018). Dietary stachyose altered the intestinal microbiota profile and improved the intestinal mucosal barrier function of juvenile turbot, *Scophthalmus maximus* L. *Aquaculture*, 486, 98-106. doi: 10.1016/j.aquaculture.2017.12.014
- Yu, L., Zzaman, W., Akanda, M. J., Yang, T., & Easa, A. (2017). Influence of superheated steam cooking on proximate, fatty acid profile, and amino acid composition of catfish (*Clarias batrachus*) fillets. *Turkish Journal of Fisheries and Aquatic Sciences*, 17, 935-943. doi: 10.4194/1303-2712-v17_5_09
- Yudiati, E., Isnansetyo, A., Murwantoko, Triyanto, & Handayani, C.R. (2019). Alginate from *Sargassum siliquosum* simultaneously stimulates innate immunity, upregulates immune genes, and enhances resistance of Pacific white shrimp (*Litopenaeus vannamei*) against white spot syndrome virus (WSSV). *Marine Biotechnology*, 21(4), 503-514. doi: 10.1007/s10126-019-09898-7