

# NATURAL RADIOACTIVITY LEVELS ALONG THE MEDITERRANEAN SAND BEACH BETWEEN TAJOURA AND MISRATA, LIBYA

Karima Elmasri and Ahmed Shetwi

Nuclear Engineering Department, Faculty of Engineering, University of Tripoli, Libya

Email: K.Elmasri@uot.edu.ly

## المخلص

تحظى قياسات مستويات النشاط الإشعاعي في الشواطئ باهتمام كبير خلال العقود الماضية لعديد الدراسات في جميع أنحاء العالم. وتوفر هذه الأبحاث امكانية رسم خرائط لمستويات الإشعاع توثيق البيانات المرجعية للتأكد من التغيرات المحتملة في النشاط الإشعاعي البيئي بسبب الأنشطة النووية والصناعية وغيرها من الأنشطة البشرية. والجدير بالذكر أن ليبيا لديها إمكانات كبيرة للسياحة مع توفر 1700 كم من السواحل.

أجريت هذه الدراسة لتحديد مستويات تركيز النشاط الإشعاعي الطبيعي في سلاسل اليورانيوم-238 والثوريوم-232 والبوتاسيوم-40 في ثلاثين عينة من رمال الشاطئ التي تم جمعها من مواقع مختلفة من الساحل بين تاجوراء ومدينة مصراته. تم تحليل عينات الرمال بواسطة نظام التحليل الطيفي لأشعة جاما باستخدام كاشف الجرمانيوم عالي النقاء (HPGe) الموجود في مختبر الإشعاع النووي في قسم الهندسة النووية بجامعة طرابلس ليبيا.

كانت تركيزات النشاط الإشعاعي لنويدات  $^{238}\text{U}$ ،  $^{232}\text{Th}$  و  $^{40}\text{K}$  بناءً على القيم المتوسطة الموزونة عبر العينات ( $0.385 \pm 6.728$ ) و ( $0.576 \pm 3.010$ ) و ( $0.183 \pm 29.673$ ) بيكريل/كجم على التوالي.

تراوح تركيز النشاط  $^{238}\text{U}$  و  $^{232}\text{Th}$  و  $^{40}\text{K}$  في عينات رمال الشاطئ من  $0.381 \pm 3.668$  إلى  $2.405 \pm 25.656$  بيكريل/كجم،  $2710 \pm 0.755$  بيكريل/كجم، إلى  $1.231 \pm 13.081$  بيكريل/كجم و  $1.460 \pm 15.935$  إلى  $6.20 \pm 88.881$  بيكريل/كجم لكل من  $^{238}\text{U}$ ،  $^{232}\text{Th}$  و  $^{40}\text{K}$  على التوالي.

تراوحت الجرعة الفعالة السنوية المكافئة في ميكروسيبرت/سنة بين  $0.494 \pm 4.092$  و  $21.694 \pm 0.214$  ميكروسيبرت/سنة بينما المتوسط العالمي للجرعة الفعالة المكافئة (AEDE) من إشعاع جاما الأرضي الخارجي أو الداخلي هو 290 ميكروسيبرت/سنة. كانت تركيزات النشاط لـ  $^{238}\text{Ra}$ ،  $^{232}\text{Th}$  و  $^{40}\text{K}$  أقل من الحد العالمي. وبالتالي لا توجد آثار إشعاعية ضارة ناتجة عن رمال البحر.

سيتم استخدام النتائج المحققة كمعلومات مرجعية لتتبع أي تغيير في مستويات الخلفية للنشاط الإشعاعي في هذه المنطقة وقد توفر أيضاً إنجازاً جيداً لرسم خرائط لخلفية النشاط الإشعاعي.

## ABSTRACT

Measurements of the levels of radioactivity in beaches are of great interest during the last decades for many studies worldwide. Such investigations can be useful for mapping of radiation levels and recording reference data to ascertain potential environmental radioactivity changes due to nuclear, industrial, and other human activities.

This study was conducted to determine the natural radioactivity concentration levels of  $^{238}\text{U}$ , and  $^{232}\text{Th}$  natural decay chains and  $^{40}\text{K}$  in 30 beach sand samples collected from different sites of the coast between Tajoura and Misrata, Libya.

Representative sand samples have been analysed based on the gamma spectroscopy system using High Purity Germanium (HPGe) detector housed in the Nuclear Radiation Laboratory at the Department of Nuclear Engineering, University of Tripoli, Libya.

The activity concentrations of  $^{238}\text{U}$ ,  $^{232}\text{Th}$  and  $^{40}\text{K}$  based on the weighted mean values across samples were  $(6.728 \pm 0.385)$ ,  $(3.010 \pm 0.576)$  and  $(29.673 \pm 0.183)$  Bq/kg, respectively.

The activity concentration of  $^{238}\text{U}$ ,  $^{232}\text{Th}$  and  $^{40}\text{K}$  in the beach sand samples ranged from  $3.668 \pm 0.381$  to  $25.656 \pm 2.405$  Bq/kg,  $0.755 \pm 0.271$  to  $13.081 \pm 1.231$  Bq/kg and  $15.935 \pm 1.460$  to  $88.881 \pm 6.20$  Bq/kg for  $^{238}\text{U}$ ,  $^{232}\text{Th}$  and  $^{40}\text{K}$  respectively. The annual effective dose equivalent in  $\mu\text{Sv/y}$  ranged between  $4.092 \pm 0.494$  and  $21.694 \pm 0.214$   $\mu\text{Sv/y}$  while the world average annual effective dose equivalent (AEDE) from outdoor or indoor terrestrial gamma radiation is  $290 \mu\text{Sv/y}$ . The activity concentrations of  $^{226}\text{Ra}$ ,  $^{232}\text{Th}$ , and  $^{40}\text{K}$  were below the global limit. Thus, there are no harmful radioactive effects resulting from sea sand.

The achieved results will be used as reference information to track any change in the radioactivity background levels in this area and may provide good achievement for mapping the radioactivity background.

**KEYWORDS:** Activity Concentration; Gamma Spectroscopy; Dose, HPGe; Activity in Beach Sand.

## INTRODUCTION

The coastline of Libya extends over 1970 km and is characterized by its relative homogeneity since there is no marked differentiation of its topography and geomorphology. Marine pollution generated by oil activities and sewage is affecting the marine environment in Libya in some areas [1]. Sand beaches produce from the rocks on land, which over time, rain, ice, wind, heat, cold, and even plants and animals break rock into smaller particles. These soil particles are transported by the flowing river to the low lying areas and deposited there as sand. The environment quality and beach areas and surroundings attract the tourists and other holidaymakers.

Environmentalists have started to put more attention to continuous monitoring of the radioactive concentration levels in the environment. Due to the serious damage that may occur to the aquatic ecosystem. There are two known main causes of the pollution; natural and anthropogenic [2, 3].

The surface contamination of water systems can be owed to increase in industrial activities and rapid urbanisation. The main source of radiation exposure for humans is the environmental natural radioactivity associated with the terrestrial and cosmic radiation. The natural radiation contributes with about 80% of a year's effective exposure for humans [4].

Individuals are exposing to radiation in two pathways; the external exposure to gamma radiation emitted from the terrestrial radionuclides;  $^{238}\text{U}$  ( $T_{1/2} = [4.468 \pm 0.003] \times 10^9$  years),  $^{232}\text{Th}$  ( $T_{1/2} = [1.405 \pm 0.006] \times 10^{10}$  years) and  $^{40}\text{K}$  ( $T_{1/2} = [1.277 \pm 0.008] \times 10^9$  years) and internal exposure from the inhalation of radioactive radon and its daughters [3-7].

The terrestrial  $\gamma$ -ray radiation with about 20% of the total annual dose (2.4 mSv) or 0.48 mSv due to natural sources and inhalation of radon as internal exposure accounting for 48% (1.15 mSv) of this dose [8-10].

Individuals long term exposure to radiation causes significant health effects. Many diseases such as lung cancers, the pancreas hepatic, due to the radon gas inhalation and ingestion of radioactive food [4,11]. Disasters, such as the Chernobyl and Fukushima, are other sources of radionuclides in the environment. Consequently, periodic radiological surveys, monitoring and screening the environment, are all necessary.

The external exposure contributes to the environmental hazardous from the natural radionuclides present in beach sands [2,5,6,12]. Measurements of radioactivity levels in beaches are of a great interest during the last decades for many studies worldwide. Such investigations can be useful for mapping of radiation levels and recording reference data to ascertain potential environmental radioactivity changes due to nuclear, industrial and other human activities [5,13,14].

Measurements of natural radioactive nuclides in beach sands have been reported in many studies worldwide. Various techniques have been used to assess natural radionuclides in different environmental media. The most common used methods include direct gamma spectroscopy, alpha spectrometry, and liquid scintillation counting [1,2,5,7,12,15-17].

The primary aim of this study is to determine the activity concentration of the natural radionuclides  $^{238}\text{U}$ ,  $^{232}\text{Th}$ , and  $^{40}\text{K}$  using gamma spectrometry based on a high purity germanium (HPGe) detector. Beach sand samples collected from the coast of Libya, the region from Tajoura to the City of Misrata, were analysed to measure and quantify natural gamma-emitting radionuclides.

## MATERIALS AND METHODS

### Sample collection and preparation

In total, 30 beach sand samples were collected from different locations on the west of Libya along the Mediterranean coast from Tajoura to the City of Misrata. Figure (1) shows these locations on the map, the distance between these sites was 4-7km. Samples of 1.5 to 2kg were collected from the closest distance from the seashore with a mold of area  $20 \times 20 \text{ cm}^2$  and a depth between 15-20 cm from the surface. Three samples were collected from each location, where the distance between them was 10 m. Samples then well-mixed and one sample was kept for each location.



**Figure 1: The regions of collected sample**

Samples were weighed, dried in oven at  $100^\circ\text{C}$  and then sieved through a 0.1-mm mesh sieve for a uniform particle size. The homogenized sample was contained in cylindrical plastic container (71 mm in diameter with 47 mm height and thickness of 1.1 mm), well-sealed with adhesive tape, weighted and labelled (according to its name, weight, and storage date). This container was chosen to match the standard source geometry.

All samples then stored for one month at the Department of Nuclear Engineering, the University of Tripoli to reach the state of the equilibrium [18,19]. Figure (2) shows the samples preparation tools and method. Table (1) lists the collected samples.



Figure 2: Samples Collection protocol

Table 1: Collected samples and their locations

Sample Label	Sample Location	Sample Label	Sample Location	Sample Label	Sample Location
Sa-1	Quara Polly	Sa-11	Ahmed Palace Resort	Sa-21	The Palm Resort (Al Khums)
Sa-2	Steam Station	Sa-12	Al Khums Port	Sa-22	Ganaima
Sa-3	Mirbat Beach	Sa-13	Zliten Port	Sa-23	Sur Sauod Beach
Sa-4	Bsees Beach	Sa-14	Zliten Dump	Sa-24	Al-Guaia HT
Sa-5	Tajoura Energy	Sa-15	Al Aman Resort	Sa-25	Toba Beach
Sa-6	Um ElZain Beach	Sa-16	Ain Kaam	Sa-26	Leptis Magna
Sa-7	Falfool Beach	Sa-17	Daphnia	Sa-27	Celine Beach
Sa-8	Trailer Factory Beach	Sa-18	Popular Market	Sa-28	Ispan
Sa-9	Iron and Steel Resort	Sa-19	Marsa Al-Quara Polly	Sa-29	Mara Tajoura
Sa-10	The Industries Resort	Sa-20	Zliten Beach	Sa-30	Zureik Beach

### Gamma-Ray Spectrometry

A high-resolution hyper-pure germanium detector-based, gamma-ray spectroscopy, low-background counting system by ORTEC was used in this study. The detector is housed in one of the nuclear radiation laboratories of the University of Tripoli at the Nuclear Engineering Department. A lead castle of approximately 10 cm thick with a copper lining on the inside to absorb any lead K X-rays, shields the detector. The relative efficiency of the detector used in these experiments is examined and compared with the manufacture nominal relative efficiency to the scintillation counter (86.4%) and found to be 76.2%. The analysis was performed using the software Ortec GammaVision® (Ortec GammaVision-32 v. 6 2010).

## Energy and Efficiency Calibration

HPGe detector energy calibration was measured over a wide range of energies. The calibration was conducted by using full-energy peaks from a standard source (SYRNORM 2005) in the shape of a cylindrical plastic container with Ra-226 activity of 9775 Bq/kg.

## Minimum Detectable Activity (MDA)

The minimum detectable activity (MDA) for each radionuclide was obtained from the background radiation spectrum for the same counting time (24hr) as for the soil samples. The values were calculated using equation (1) [20].

$$MDA = \frac{N_d}{\varepsilon P_\gamma T m} \quad (1)$$

Where;  $N_d = L_c + 2.706$ ,  $L_c = 4.653 \sigma_{NB}$

Where,  $N_d$ ,  $\sigma_{NB}$ ,  $\varepsilon$ ,  $P_\gamma$ ,  $L_c$  are the counts, the standard deviation of the background, detector efficiency, the emission probability of the gamma line corresponding to the peak energy, critical level, respectively, and T is the sample counting time.  $m$  is sample mass.

## RESULTS AND DISCUSSION

### Detection Limit and Minimum Detectable Activity (MDA)

The minimum detectable activity obtained was in the range of  $0.110 \pm 0.001$  to  $1.01 \pm 0.002$  Bq/kg as shown in Figure (3). This range of values was used to check if the activity concentrations found within the samples are of measurable values throughout this study.




Figure 3: Calculated MDA for listed radionuclides

### Activity Concentrations

The activity concentrations of each identified radionuclide from  $^{238}\text{U}$ ,  $^{232}\text{Th}$  series and  $^{40}\text{K}$  of seashore sand samples were calculated using the following equation:-

$$A(\text{Bq/kg}) = \frac{N-B}{\varepsilon \times t \times I_\gamma \times m} \quad (2)$$


Where  $\varepsilon$  is the absolute efficiency at a given photo peak energy,  $I_\gamma$  is the emission probability of the gamma line corresponding to the peak energy,  $m$  is the mass of the measured sample (kg),  $N$  and  $B$  are the areas under the photopeaks of the sample and the background, respectively [5,12].

The gamma-energy lines of 609.32 (45.4%) keV, 1120.29 (14.8%) keV, emitted by  $^{214}\text{Bi}$  were used to represent the  $^{238}\text{U}$  ( $^{226}\text{Ra}$ ) series, while 911.20 (25.8%) keV and 968.97 (15.8%) keV from  $^{228}\text{Ac}$  were used to represent the  $^{232}\text{Th}$  series and 1460.82 (10.66%) keV for  $^{40}\text{K}$  gamma-line.

Then obtained values were compared with the reported natural radionuclides mean activities in Libyan soil by UNSCEAR (2008) global survey on exposures to natural radiation sources of  $^{238}\text{U}$ ,  $^{232}\text{Th}$  series and  $^{40}\text{K}$ , which are 10.5, 8.8, 9.5 and 270 Bq/kg, respectively [20].

In addition, these activities were compared with world limit (33 Bq/kg for  $^{238}\text{U}$ , 45 Bq/kg for  $^{232}\text{Th}$ , and 420 Bq/kg for  $^{40}\text{K}$ ) reported by UNSCEAR 2008. The activity concentration of  $^{238}\text{U}$ ,  $^{232}\text{Th}$  and  $^{40}\text{K}$  based on the weighted mean values across samples as shown in Figures (4-6) were  $(6.728 \pm 0.385)$ ,  $(3.010 \pm 0.576)$  and  $(29.673 \pm 0.183)$  Bq/kg, respectively. The lowest radioactivity of  $^{238}\text{U}$  was about  $(3.668 \pm 0.381)$  Bq/kg in sand sample Sa-01 collected from Quara Polly. Sample Sa-9 from Iron and Steel Resort contained the lowest activity of  $^{232}\text{Th}$ ,  $(0.755 \pm 0.271)$  Bq/kg.

For  $^{40}\text{K}$  the lowest activity concentration obtained in sample Sa-22 collected from Ganaima  $(15.935 \pm 1.460)$  Bq/kg, while the largest activity recorded in sample S-14 from Zliten Dump with  $(88.881 \pm 6.201)$  Bq/kg). Sample S-20 collected from Zliten Beach  $(32^\circ 30' 21.1''\text{N } 14^\circ 30' 01.6''\text{E})$  revealed the highest values of  $^{238}\text{U}$  and  $^{232}\text{Th}$  series  $(25.656 \pm 2.405)$  and  $(13.081 \pm 1.231)$  Bq/kg respectively. These values were higher than limits of activity of natural radionuclides activity in Libyan soil reported in UNSCEAR (2008).



**Figure 4: Ra-226 radioactivity concentrations in Bq/kg.**




Figure 5: Th-232 radioactivity concentrations in Bq/kg




Figure 6: K-40 radioactivity concentrations in Bq/kg

## Radiation Hazard Assessment

### Gamma Absorbed Dose Rate Assessment

The Radium equivalent activity in each sample was calculated by:

$$Ra_{eq} = A_{Ra} + 1.43A_{Th} + 0.077A_K \quad (3)$$

Where  $A_{Ra}$ ,  $A_K$  and  $A_{Th}$  are the specific activities of  $^{238}U$ ,  $^{40}K$  and  $^{232}Th$  in (Bq/kg).

The values of  $Ra_{eq}$  found to be in the range of  $7.743 \pm 0.359$  to  $44.681 \pm 0.149$  and Bq/kg, which were below the world's mean values 370 Bq/kg [14,16,21].

The gamma dose rate (D) in nGy/hr caused by naturally occurring radioactive materials in air at 1 m above the ground surface can be estimated using the following formula [5]:-

$$D = 0.461A_{Ra} + 0.623A_{Th} + 0.0414A_K \quad (4)$$

Where  $A_{Ra}$ ,  $A_{Th}$  and  $A_K$  are the specific activities of  $^{226}Ra$ ,  $^{232}Th$  and  $^{40}K$  in (Bq/Kg).

The range of calculated gamma dose rate in this study was found to be between  $4.092 \pm 0.494$  and  $21.69 \pm 0.214$  nGy/h. Figure (7) illustrates the Dose rate,  $Ra_{eq}$  distribution for all samples.




Figure 7: Dose rate,  $R_{aeq}$  distribution for all samples.

### External Hazard and Annual effective dose equivalent (AEDE)

External hazard index for each sample was found to be less than unity, and Table (2) lists these results, which calculated by:-

$$H_{ex} = \frac{C_K}{4810} + \frac{C_{Th}}{259} + \frac{C_{Ra}}{370} \quad (5)$$

Where  $C_K$ ,  $C_{Th}$  and  $C_{Ra}$  are the activities concentrations of  $^{40}\text{K}$ ,  $^{232}\text{Th}$  and  $^{226}\text{Ra}$  (in Bq/kg) respectively. This expression indicates that the value of this index must be less than unity for the radiation hazard to be insignificant. Thus, the maximum values of  $H_{ex}$  equal to unity correspond to the upper limit of  $R_{aeq}$  being 370 Bq/kg.

Table 2: External Hazard Index from 30 Samples.

Sample #	$H_{ex}$	Sample #	$H_{ex}$	Sample #	$H_{ex}$
Sa-1	0.032	Sa-11	0.028	Sa-21	0.029
Sa-2	0.032	Sa-12	0.032	Sa-22	0.023
Sa-3	0.035	Sa-13	0.030	Sa-23	0.030
Sa-4	0.048	Sa-14	0.041	Sa-24	0.034
Sa-5	0.035	Sa-15	0.037	Sa-25	0.030
Sa-6	0.028	Sa-16	0.028	Sa-26	0.031
Sa-7	0.028	Sa-17	0.036	Sa-27	0.027
Sa-8	0.031	Sa-18	0.046	Sa-28	0.038
Sa-9	0.036	Sa-19	0.039	Sa-29	0.051
Sa-10	0.030	Sa-20	0.128	Sa-30	0.034


### The Annual Effective Dose Rate

In order to estimate the annual effective dose rate in air, the conversion coefficient from absorbed dose in air to effective dose received by an adult must be considered. This value is published in UNSCEAR 2000 and UNSCEAR 1993, to be  $0.7 \text{ SvGy}^{-1}$  for environmental exposure to gamma rays of moderate energy. The outdoor occupancy factor is about 0.2. The annual effective dose equivalent is given by the following equation [14].

$$\text{AEDE } (\mu\text{Sv/y}) = D(\text{nGy/h}) \times 8760(\text{h/y}) \times 0.2 \times 0.7(\text{Sv/Gy}) \times 10^{-3} \quad (6)$$

The annual effective dose equivalent in  $\mu\text{Sv/y}$  shown in Figure (8) ranged between 5.019 and 26.606  $\mu\text{Sv/y}$ , which was within the allowed limits that equal (1 mSv/y) for all samples and all regions that were selected in the present study.





**Figure 8: annual effective dose equivalent for each sample**

## CONCLUSION

Gamma spectrometer of HPGe was used to measure the activity concentration of natural terrestrial radionuclides,  $^{238}\text{U}$ ,  $^{232}\text{Th}$ , and  $^{40}\text{K}$  in beach sands collected from different sites of the coast between Tajoura and Misrata, Libya. Reported values are comparable to the average worldwide range.

The achieved results were compared with previously published study, which was conducted to measure radioactivity concentrations of beach sand samples from different locations along the coast of Tripoli, northwest Libya, using high resolution  $\gamma$ -ray spectroscopy. Their findings showed that the radioactivity concentrations for  $^{226}\text{Ra}$ ,  $^{232}\text{Th}$ ,  $^{40}\text{K}$  in samples taken at depth 5-10 cm have an average  $7.5 \pm 2.5$  Bq/kg,  $4.5 \pm 1.3$  Bq/kg,  $28.5 \pm 6.7$  Bq/kg respectively. This study showed that the activity of  $^{238}\text{U}$ ,  $^{232}\text{Th}$  and  $^{40}\text{K}$  based on the weighted mean values across samples were  $(6.728 \pm 0.385)$ ,  $(3.010 \pm 0.576)$  and  $(29.673 \pm 0.183)$  Bq/kg, respectively, which is consistent with the previously published study [21].

The range of the absorbed dose rate obtained for the soil samples was between 2.7n Gy/h and 6.1 nGy/h, while the mean value of the absorbed dose rates in air was  $4.4 \pm 1.3$  nGy/h. In this work the absorbed dose rate values were higher and found to be between  $4.092 \pm 0.494$  and  $21.69 \pm 0.214$  nG/h. [21].

All the results obtained in this study showed that the beaches in the region of study are safe in terms of radiological hazard, with no estimated values exceeding the permissible levels. Obtained results will be employed as reference information to track any change in the radioactivity background levels within the area covered in this study and may provide good achievement for mapping the radioactivity background.

## REFERENCES

- [1] "National Integrated Monitoring and Assessment Programme (Imap) for Coast and Hydrography Indicators for Libya," 2013.
- [2] S. Chandrasekaran, G. Sankaran Pillai, and B. Venkatraman, (2020), "Assessment of heavy metals and radionuclides ( $^{238}\text{U}$ ,  $^{232}\text{Th}$  and  $^{40}\text{K}$ ) concentration of beach sands collected from East Coast of Tamilnadu, India with multivariate statistical approach", *Int. J. Environ. Anal. Chem.*, pp. 1–23. (Volume 102, 2022 - Issue 16)
- [3] A. E. Abdel Gawad *et al.*, (2021), "Radiological investigation on sediments: A case study of Wadi Rod Elsayalla the southeastern desert of Egypt," *Appl. Sci.*, vol. 11, no. 24.
- [4] UNSCEAR, 2000. Exposures from natural radiation sources (Annex B). Sources and Effects of Ionizing Radiation. UNSCEAR: (2000); New York, NY, USA 84–141.

- [5] F. Caridi *et al.*, (2016). "Natural radioactivity and elemental composition of beach sands in the Calabria region, south of Italy," *Environ. Earth Sci.*, vol. 75, no. 7, 2.
- [6] UNSCEAR, (2010). *Scientific Report: summary of low-dose radiation effects on health*.
- [7] M. Awad, et.al, (2022) "Radioactive risk assessment of beach sand along the coastline of Mediterranean Sea at El-Arish area, North Sinai, Egypt", *Mar. Pollut. Bull.*, vol. 177, p. 113494,.
- [8] IAEA, (2003). "Radiation protection and the management of radioactive waste in the oil and gas industry," *Environ. Health Perspect.*, vol. 34, no. 34, p. 130.
- [9] IAEA. (2003), "Extent of environmental improvement", *Int. At. Energy Agency Aust.*, vol. 24, no. 4, pp. 433–434.
- [10] J. E. Ndjana Nkoulou II, et al., (2018). "Natural Radioactivity Measurements in Soil, External Dose and Radiological Hazard Assessment in the Uranium and Thorium Bearing Region of Lolodorf, Cameroon," *Radioisotopes*, vol. 67, no. 9, pp. 435–446, 2018.
- [11] C. Portier, (2003 ). "Draft Toxicological Profile for Radon: Agency for Toxic Substances and Disease Registry (ATSDR)," *U.S. Dep. Heal. Hum. Serv.*, September, p. 283.
- [12] P. Sola, U. Injarean, et al., (2022). "Naturally occurring radioactive materials in sea water and beach sand near Map Ta Phut Industrial Estate, Thailand," *Maejo Int. J. Sci. Technol.*, vol. 16, no. 2, pp. 98–111.
- [13] M. M. Orosun, et al., (2019). "Natural radionuclides and radiological risk assessment of granite mining field in Asa, North-central Nigeria," *MethodsX*, vol. 6, November, pp. 2504–2514.
- [14] M. T. S. Heikal, et al., (2017). "Environmental assessment of radiation impacts on Sharm El-Sheikh Area, South Sinai, Egypt," *J. King Abdulaziz Univ. Earth Sci.*, vol. 24, no. 1, pp. 57–78
- [15] S. Fares, (2017). "Measurements of natural radioactivity level in black sand and sediment samples of the Tamsah Lake beach in Suez Canal region in Egypt," *Appl. Sci.*, vol. 10, no. 3, pp. 194–203.
- [16] H. S. Eissa, et al., (2012). "Radiation dose estimation of sand samples collected from different Egyptian beaches," *Radiat. Prot. Dosimetry*, vol. 147, no. 4, pp. 533–540.
- [17] V. Tech Artūras Jukna, and Gražina Grigaliūnaitė-Vonsevičienė (2022). "Assessing the Radioactivity Level of Seawater and Sand Samples from the Baltic Sea Region", Research Square article 2022. pp. 1–14.
- [18] R. W. D. Garcêz, et al., (2019). "Study of K-40, Ra-226, Ra-228 and Ra-224 activity concentrations in some seasoning and nuts obtained in Rio De Janeiro city, Brazil," *Food Sci. Technology.*, vol. 39, no. 1, pp. 120–126.
- [19] U. Elghawi, et al., (2019). "Studying of Naturally Occurring Radioactive Materials (NORM) in Oilfield (A/100) South East of Libya. Environmental Research Journal
- [20] Kohn, Henry I. (1989) "Sources, effects and risks of ionizing radiation." *Radiation Research* 120.1: 187.
- [21] S. U. El-Kameesy, et al., (2008). "Natural radioactivity of beach sand samples in the Tripoli Region, northwest Libya," *Turkish J. Eng. Environ. Sci.*, vol. 32, no. 4, pp. 245–251.