



Fertility and Reproductive Outcome in Mice Following Trichloroethane (TCE) Exposure

M. A. Al-Griw¹, S. A. Al-Azreg², E. M. Bennour³, S. A. M. El-Mahgiubi², A. R. Al-Attar², N. M. Salama¹, A. Elnfati¹

1. Developmental Biology Division, Zoology Department, Faculty of Science, University of Tripoli, Tripoli-Libya
2. Department of Pathology and Clinical Pathology, Faculty of Veterinary Medicine, University of Tripoli, Tripoli-Libya
3. Department of Internal Medicine, Faculty of Veterinary Medicine, University of Tripoli, Tripoli-Libya.

*Corresponding author: Abdul Hakim Elnfati

Abstract: Exposure to trichloroethane (TCE), an industrial solvent, has been shown to be negatively associated with reproductive performance. The present study was performed to assess the effects of TCE exposure on the reproductive performance and outcome in mice during a critical developmental window of later reproductive life. A group of female mice were injected intraperitoneally twice weekly for three weeks with TCE (100 and 400 µg/kg). Mice were followed up for signs of toxicity and death. Changes in uterine tissues have also been investigated by histopathology. The results showed that TCE exposure has reduced the number of F0 fertile females comparing to controls. Moreover, TCE exposure resulted in a decreased pups number and changed sex ratio in the litter of F0 TCE-treated dams. Histopathological examination revealed a TCE-induced uterine toxicity appeared as a severe endometrial hyperplasia with squamous cell metaplasia and adenomyosis. These results indicate that TCE exposure during a critical reproductive developmental window could affect the fertility and interfere with the reproductive outcome in mice.

Keyword: TCE; fertility; reproductive outcome; uterus; mice.

INTRODUCTION

Infertility is a major worldwide public health concern because it affects approximately 10% of all reproductive-aged couples¹. Exposure to chemicals during reproductive developmental windows may predispose individuals to disease and/or dysfunction later in life. 1,1,1-trichloroethane (TCE), an ambiguous environmental pollutant, is widely used as an industrial solvent and a degreasing agent²⁻⁴. It has been shown that TCE is well absorbed by all routes of exposure. At least by inhalation, the rate of uptake is driven initially by tissue loading and then by metabolism once steady state conditions have been reached¹. Numerous studies with developmental exposure to lower doses than the “safe” dose suggest that TCE exposure causes various detrimental defects, such as low fetal weight, birth defects, and developmental disorders. Recent epidemiological studies have shown that chemical exposure environmentally or occupationally on a daily basis is associated with increase a woman's risk of spontaneous abortions, infertility, low fetal weights, birth defects¹.

However, to the best of the author's knowledge, it has not been examined whether TCE exposure, during a critical uterine developmental window, has consequences on reproductive performance (e.g. fertility, gestation and litter life) in later reproductive life. Thus, this study was performed to investigate the long-term effects of environmentally relevant low and high dose of TCE on adult reproductive functions, such as fertility and reproductive outcome.

MATERIAL AND METHODS

Animals and housing

A total of thirty six female Swiss Albino mice (F0 generation), with an age range of 4-6 weeks and weight range of 21–24 g, were used in this study. Mice were kept under a constant light-dark cycle (dark period from 7:00 pm to 7:00 am) at $24 \pm 1^\circ\text{C}$ and $55 \pm 5\%$ relative humidity. Food and drinking water were available ad libitum. All efforts were made to minimize the pain during animal handling and experimentation and to reduce the number of animals used.

Study design and treatment regimen

TCE (Baxter International) was suspended in corn oil. Female mice (F0) were divided into four groups of six mice each. These groups were: the low and high dose TCE-treated (100 or 400 $\mu\text{g}/\text{kg}$) groups and the vehicle and sham control groups. For TCE-treated groups, the doses were calculated and delivered in 80-100 μl of corn oil based on their body weight⁵⁻⁶. TCE doses were selected as they considered safe by Environmental Protection Agency (EPA)⁷. Vehicle controls were received an equal volume of corn oil only. The sham controls were not received any exposure. TCE or vehicle were administered intraperitoneally at a defined time (10:00 am) every 3rd day. The exposure window was selected because this is the critical development window in the mouse¹.

After treatment, F0 female mice were mated with fertility confirmed control males (2 females:1 male ratio). Mating was confirmed by the presence of vaginal plug. Once the plug was observed, females were separated from males and individually caged. The day the vaginal plug detected was defined as the first gestation day (GD1). Dams (F0 generation) were observed daily and body weight gain was measured daily to further confirm pregnancy and for any adverse clinical signs or abnormal behavior that may result from toxicity. The dams were allowed to deliver naturally and the delivery day was designed as postnatal day 0 (PND0).

The study comprehensive teratological parameters for F0 females included body weight, reproductive performance (fertility index; gestation index, mortality), and gross pathology. The study parameters for F1 offspring included pup size, average pup weight, sex ratio, pup mortality and stillbirths as recorded on PND0.

After the two-week mating period, unmated females were singly housed, observed for estrous cycle for another 10 days, and body weight was monitored for another two weeks. The females were considered infertile if they did not cycle and/or did not have significant body weight gain during the entire testing period.

Clinical assessment

The clinical assessment included animal survival, body weight gain and histopathology.

Animal survival

During the course of the exposure period, mice were observed twice per day for any abnormal clinical signs or behavior that may result from toxicity. Mice were assessed for morbidity and mortality twice daily, midmorning and late afternoon. Night deaths were recorded the next morning. Two independent observers confirmed the cause of death to exclude TCE-nonrelated mortality.

Body weight

Mouse body weight in control and TCE-treated groups was assessed on a weekly basis to monitor the effect of TCE exposure on body weight.

Histopathological examination

After dissection, 10%-formalin-fixed uteri were processed in a series of graded ethanol solutions and embedded in paraffin wax. Paraffin sections were cut at 6-8 μm thickness, deparaffinized, rehydrated, stained with hematoxylin and eosin (H&E) and examined under a light microscope (Leica, Germany) for histopathology.

Fertility and Reproductive Outcome in Mice Following ...

Statistics

Data were expressed as means \pm SEM (standard error of the means) from 6 female mice of each group using SPSS software, version 20. A computerized Kolmogorov-Smirnov test was used to determine whether the data fitted a normal distribution. One-way ANOVA test followed by Tukey's post hoc comparisons was used to make multiple comparisons between treatment groups. Student's t-tests were used to make comparisons between two groups. Mann-Whitney U-test was used for nonparametric samples. Statistical significance was assigned at $P \leq 0.05$.

RESULTS

Effect of TCE on animal survival

No mortality has been recorded among mice in all groups along the course of the experiment except for one death case out of the six females in 400 $\mu\text{g}/\text{kg}$ TCE-treated group recorded four weeks post TCE exposure.

Effect of TCE on body weight

Previous animal and epidemiological studies have linked developmental chemical/ toxicant exposure to metabolic disorder and obesity¹. Therefore, in this study we decided to investigate the effect of TCE exposure on body weight of F0 female mice by monitoring the body weight either pre- and post-treatment.

In this study, TCE exposure have not shown an effect on overall body weight in F0 females as statistical analysis indicated no significant between treated and non-treated groups (data not shown). However weight gain in F0 females in 400 $\mu\text{g}/\text{kg}$ TCE-treated group, but not 100 $\mu\text{g}/\text{kg}$ TCE-treated group ($P = 0.332$) was significantly higher ($P = 0.024$) comparing to controls (Figure1). This concludes that TCE exposure at the dose of 100 $\mu\text{g}/\text{kg}$ is a No Observed Adverse Effective Level (NOAEL) for F0 female mice in this study.




Figure 1. Body weight gain of the F0 females. TCE exposure significantly increased weight gain at a dose of 400 $\mu\text{g}/\text{kg}$. Data represent mean \pm SEM of $n = 6$ animals per groups. # Significantly different from the vehicle controls ($P \leq 0.05$).

Effect of TCE on fertility

To determine the effect of TCE exposure on fertility in later reproductive life and whether the effects would change with age, a number of fertility and reproductive indicators including percentage of fertile female has been investigated.

The results showed an effect of TCE exposure on the fertility of F0 females. The analysis showed a reduction in the percent of fertile females in TCE-treated groups as in 100 $\mu\text{g}/\text{kg}$ TCE-treated group, five females out of six females (~83%) gave birth while in 400 $\mu\text{g}/\text{kg}$ TCE-treated group, only one female out of six females (~17%) gave birth (Figure 2).

Statistically, the fertility was significantly reduced in 400 $\mu\text{g}/\text{kg}$ TCE-treated F0 females comparing to controls ($P = 0.047$; Figure 2).




Figure 2. Effect of TCE exposure on fertility of F0 females. The fertility was significantly reduced in 400 $\mu\text{g}/\text{kg}$ TCE-treated F0 females. Data represent mean \pm SEM of $n = 6$ animals per groups. # Significantly different from the vehicle controls ($P \leq 0.05$).

Effect of TCE on reproductive outcome

To determine the effect of TCE exposure on reproductive outcome, a number of reproductive indicators including litter number and mortality, pup size and weight and sex ratio have been investigated. The results showed also that TCE exposure at a dose of 100 or 400 $\mu\text{g}/\text{kg}$ significantly affected the litter number and decreased the average number of pups, especially for the TCE 400 $\mu\text{g}/\text{kg}$ TCE-treated group compared to controls (Figure 3A). However, TCE exposure at a dose of 100 or 400 $\mu\text{g}/\text{kg}$ had no effect on pup size comparing to controls (data not shown).

For the effect of TCE exposure on average live pup weight, TCE exposure at a dose of 100 $\mu\text{g}/\text{kg}$ had no effect on the pup weight ($P = 0.388$) comparing to controls. However, TCE exposure at a dose of 400 $\mu\text{g}/\text{kg}$ has significantly ($P = 0.049$) increased the average of live pups weight comparing to controls (Figure 3B). Furthermore, the results of this study showed that TCE exposure at either a dose of 100 or 400 $\mu\text{g}/\text{kg}$ had no effect on litter mortality when compared with controls.

Fertility and Reproductive Outcome in Mice Following ...




Figure 3. Effect of TCE exposure on reproductive life/ outcome. (A) Quantification of pup number. (B) Measurement of pup weight. Data represent mean \pm SEM of $n = 6$ animals per groups. # Significantly different from the vehicle controls ($P \leq 0.05$).

Regarding the effect of TCE exposure on the litter sex ratios, TCE exposure had a significant effect on the male/female ratio at a dose of 400 µg/kg. This dose has significantly changed the litter sex ratio in favor of F1 males comparing to controls ($P < 0.05$) (Figure 4). However, neither 100 µg/kg TCE dose exposure nor the vehicle had effect on the litter sex ratio (Figure 4).




Figure 4. Effect of TCE exposure on reproductive outcome. TCE exposure at a dose of 400 µg/kg has significantly changed the litter sex ratio in favor of F1 males comparing to controls. Data represent mean \pm SEM of $n = 6$ animals per groups. # Significantly different from the vehicle controls ($P \leq 0.05$).

Gross pathological findings and histopathological changes of uterine tissues

At necropsy, no gross pathological changes have been noticed on control mice. In addition, no histopathological changes were observed in their uterus (Figure 5A) or the other parts of reproductive tract. However, abilateral asymmetrical enlargement was seen in the uterine body of several TCE-treated mice at a dose of 100 and 400 µg/kg. Sections from different parts of uterus were submitted to histopathological examination.

The histopathological examination of uterine tissues of TCE-treated mice at a dose of 100 µg/kg showed a marked increase in the endometrial and myometrial thickness (Figure 5B). This thickening is revealed as diffuse or focal proliferative reactions with characteristic hyperplasia of uterine glands with prominent stromal elements and/or formation of intra-luminal polyp. The glands were large, irregular, highly branched and/or cystic lined by single or double layer of non-ciliated secretory epithelium (Figure 5C). In some cases, polyps were appeared as evaginated circumscribed mass covered by columnar epithelium with underlying tissue formed from endometrial stromal elements (Figure 5D). The myometrium showed normal smooth muscles fiber and vascular constituent, however, some cases showed invaginated uterine glands in-between (Adenomyosis) (Figure 5E). These histopathological changes caused by TCE were remarkably increased in mice which were treated with 400 µg/kg TCE; the endometrial glands were severely ectatic, lined by flattened epithelial cells and contain few degenerative heterophils and necrotic cellular debris (Figure 5F) and most endometrial polyps were composed of abundant amounts of loosely or compactly arranged spindle-shaped or satellite endometrial glands (Figure 5G). Moreover, there was sever hyperplastic changes with squamous cell metaplasia in endometrial epithelium (Figure 5H). In conclusion, the uteri of TCE-treated mice showed a severe endometrial hyperplasia with squamous cell metaplasia and adenomyosis.




Figure 5. Representative uterine sections of control and TCE-treated mice (H&E staining). A. Uterine sections of control mice showing normal tissue architecture at magnification X200. B. An endometrial thickening with more $\mu\text{g}/\text{kg}$ TCE-treated mice. C. TCE-treated mice showing prominent hyperplastic uterine glands and stromal elements in endometrium showing hyperplastic glands (cystic endometrial hyperplasia) (asterisk). D. Pedunculated endometrial polyp with an epithelial covering and stromal hyperplasia in $100 \mu\text{g}/\text{kg}$ TCE-treated mice (asterisk). E. Endometrial tissue includes glands and stromal elements within the myometrium (arrows) in $100 \mu\text{g}/\text{kg}$ TCE-treated mice. F. The endometrial glands are severely ectatic, lined by flattened epithelial cells and contain few degenerative heterophils, necrotic cellular debris and secretory material (asterisk). G. Papillary fibroepithelial polyp arising from the endometrium with cystic change of uterine glands in $400 \mu\text{g}/\text{kg}$ TCE-treated mice.

(asterisk). H. Hyperplasia and squamous metaplasia of epithelium of uterine glands in 400 µg/kg TCE-treated mice (arrows).

DISCUSSION

The exposure to environmental toxic chemicals causes many harm effects in biological cell systems^{2-3,5,8-10}. TCE is a non-carcinogenic (group 3) because there is inadequate evidence for carcinogenicity in both human and animals according to the last update of U.S Environmental Protection Agency (EPA) and National Toxicology Program (NTP) technical report¹¹⁻¹². In addition, according to WHO toxicological report, TCE is not considered as toxic, and not necessary to drive a health based guideline standard¹³. However, the Agency for Toxic Substances and Disease Registry (ATSDR) indicated that TCE affects many internal organs, such as cardiovascular and nervous system¹⁴.

Despite the TCE safe profile claimed in the aforementioned reports, TCE reproductive toxicity was reported by several studies in many animal models by oral, inhalation and dermal exposure¹⁵⁻¹⁸. Moreover, prior animal studies have shown that perinatal exposure to TCE affects the development of the brain, liver, adipose tissue and reproductive tract and adversely affects their functions^{2-3,5,8-10}. However, based on the best authors knowledge, no study had focused on the long-term effect of TCE exposure on the fertility in later reproductive life.

Thus, this study was designed to explore the possible hazard effects of the exposure to environmentally relevant levels of TCE during an embryonic ovarian developmental window has long lasting effects on reproductive performance in later reproductive life. The administration of TCE safe considered doses to female mice was through intra-peritoneal injection. This mode of exposure has never been reported previously but the advantage of such exposure is to put the toxic chemical in close contact with the target cells and avoid rapid bio-elimination of TCE, which is two hours¹⁹.

In the present study, TCE exposure impaired the fertility and increased body weight gain in F0 females. In addition, most interestingly, TCE exposure also affected the reproductive outcome of F0 females.

The current data showed also that TCE had not resulted in a change in the body weight of F0 female mice. This is consistent with a previous study on rats which reported that no body weight changes were observed following intermittent or continuous exposure to trichloroethylene vapors at exposure levels in a range of 400–2,500 ppm for 2–13 weeks²⁰. On the other hand, another study reported > 20% decreased body weight in male rats exposed to trichloroethylene vapors at 376 ppm for 4 hours per day, 5 days per week and a period of 12 or 24 weeks²¹. These findings are not consistent with other studies, which have shown that developmental TCE exposure disturbs metabolism, interferes with adipocyte proliferation and differentiation and may be linked to obesity in later life²⁰⁻²¹. However, as no significant differences in body weight have been observed between TCE-treated mice and controls, we speculate in TCE exposure might reprogram the progenitor adipocytes and increase body weight significantly later in life.

Although some studies indicated that the exposure to TCE decreases body weight gain¹⁷, the results of this study showed a dose-associated increase in the body weight gain of TCE-treated F0 females and their neonates. Many studies have found that certain toxic chemicals such as pesticides, the most environmental pollutants, and plastic products can cause weight gain by disrupting endocrine system²²⁻²⁶. These environmental chemicals have different mechanism of action such as increasing the activity of estrogen^{23,25}, abnormal adipogenesis²⁵, increasing inflammatory cytokine activity, causing oxidative stress, inducing abnormal thyroid function and impacting energy metabolism²⁷. In addition, inspite of rare information of TCE endocrine toxicity, one study recorded that acute inhalation of TCE disrupts the concentration of corticosterone and adrenocorticotrophic hormones²⁸. It was also noted that TCE has increased the weight of internal organs of rats²⁹ and their pups³⁰⁻³¹. In this study, it is possible that the

Fertility and Reproductive Outcome in Mice Following ...

increase in body weight gain due to TCE exposure has been caused by one or more of above mentioned mechanisms.

Many toxicological and histopathological examinations in different animal models indicated that exposure to TCE through inhalation has not adverse effects on female reproductive system following wide range doses (acute, intermediate and chronic) ^{15,29,32-35}. Similar to TCE inhalation, oral exposure of 100, 300 and 1000 mg/kg/day in mice [36] and 3 mg/kg in rats ³⁷⁻³⁸ did not show any hazard effects on female body weight, survival, fertility, reproductive performance and gestation period. In this study, our model (with low doses) showed a significant decrease in the pub number in TCE-treated female litter. This might be attributed to the effect of TCE on oogenesis and oocyte quality, fertilization processes, early embryo death or the changes noticed in endometrium which could affect the proper formation and implantation of placenta.

Abnormal sex ratio may be induced in human and animals by the exposure to chemicals ³⁹⁻⁴⁰ but changes in sex ratio related to exposure to TCE have not been documented. This study is providing the first observation about changed litter sex ratio in mice due to exposure to high doses of TCE. This could be attributed to an increased sensitivity of sperms and/or embryos of certain genotype to the exposure to a particular chemical resulting in low fertilization rate or early embryonic death ⁴¹.

The uterus is one of the major organs of reproductive activity of mammals as it provides the place and the environment for gestation from the embryonic implantation and placentation till parturition. Therefore, the pathological conditions interfering with the normal uterine physiology might result in impaired reproductive performance. This study showed that TCE exposure induced a bilateral asymmetrical enlargement in the uterine body. Histopathologically, TCE exposure was associated with endometrial tissue changes appeared as severe endometrial hyperplasia and adenomyosis. This might be the responsible, at least in part, to the reduced fertility in TCE-treated F0 females and their reduced litter number.

In conclusion, the current study indicates that TCE exposure during a critical ovarian developmental window impairs female reproduction life. Further studies focused on examining the hormone profiles, ovarian morphology at later reproductive life, and pregnancy status during mid- and late gestation will be helpful in enhancing our understanding of the mechanism by which TCE affects female reproduction.

Acknowledgements

This investigation was supported in part by the Division of Developmental Biology, Faculty of Science, University of Tripoli, Tripoli-Libya. The authors would like to thank Turkia Aduma, the technical staff at the Department of Pathology and Clinical Pathology, Faculty of Veterinary Medicine, University of Tripoli, Tripoli-Libya, for her technical assistance.

REFERENCES

1. Wang W, Hafner KS, Flaws JA. In utero bisphenol A exposure disrupts germ cell nest breakdown and reduces fertility with age in the mouse. *Toxicology and Applied Pharmacology*. 2014; 276:157-164.
2. House RA, Liss GM, Wills MC, Holness DL. Paresthesias and sensory neuropathy due to 1,1,1-trichloroethane. *J. occup. environ. Med*, 1996; 38: 123-124.
3. Wang G, et al., Oxidative and nitrosative stress in trichloroethene-mediated autoimmune response. *Toxicology*. 2007; 229: 186–193.
4. York RG, Sowry BM, Hastings L, Manson JM. Evaluation of teratogenicity and neurotoxicity with maternal inhalation exposure to methyl chloroform. *Journal of Toxicol. environ. Health*. 1982; 9: 251-266.
5. Wang, G., et al., N-Acetylcysteine protects against trichloroethene-mediated autoimmunity by attenuating oxidative stress. *Toxicology and Applied Pharmacology*. 2013. 273: p. 189-195.

6. Melani, A., et al., The selective A2A receptor antagonist SCH 58261 reduces striatal transmitter outflow, turning behavior and ischemic brain damage induced by permanent focal ischemia in the rat. *Brain Research*. 2003; 959: 243-250.
7. Lane RW, Riddle BL, Borzelleca JF. Effects of 1,2-dichloroethane and 1,1,1-trichloroethane in drinking water on reproduction and development in mice. *Toxicol Appl Pharmacology*, 1982; 63: 409–421.
8. Topham JC. Do induced sperm-head abnormalities in mice specifically identify mammalian mutagens rather than carcinogens. *Mutat Res*. 1980; 74: 379-387.
9. Griffin JM, Blossom SJ, Jackson SK, Gilbert KM, Pumford NR. Trichloroethylene accelerates an autoimmune response by Th1 T cell activation in MRL+/- mice. *Immunopharmacology*. 2000; 46: 123-137.
10. Snyder R, Andrews LS. Toxic effects of solvents and vapors. In: Klaassen, CD; ed. Casarett and Doull's Toxicology: The Basis Science of Poisons. 5th ed. New York: McGraw-Hill, 1996.
11. NTP. NTP technical report on the toxicity studies of 1,1,1-trichloroethane administered in microcapsules in feed to F344/N rats and B6C3F1 mice. National Toxicology Program. (41) NIH 004402. 2000.
12. EPA. Toxicological Review of 1,1,1-Trichloroethane (CAS No. 71-55-6) In Support of Summary Information on the Integrated Risk Information System (IRIS). . EPA/635/R-03/013. U.S. Environmental Protection Agency, Washington, DC. 2007.
13. WHO. 1,1,1-Trichloroethane in Drinking-water. Background document for development of WHO Guidelines for Drinking-water Quality. World Health Organization. 2003; WHO/SDE/WSH/03.04/65.: p. 16.
14. ASTDR. 1,1,1 Trichloroethane. Agency for Toxic Substances and Disease Registry. 2006; p. 371.
15. Quast J, Calhoun LL, and Frauson LE. 1,1,1-trichloroethane formulation: a chronic inhalation toxicity and oncogenicity study in Fischer 344 rats and B6c3F1 mice. *Fundamental and applied toxicology* . 1988; 11(4): 611-25.
16. Tyson CA, et al., Correlations of in vitro and in vivo hepatotoxicity for five haloalkanes. *Toxicology and applied pharmacology*. 1983; 70(2): 289-302.
17. Bruckner JV, et al., Acute, short-term, and subchronic oral toxicity of 1,1,1-trichloroethane in rats. *Toxicological sciences : an official journal of the Society of Toxicology*. 2001, 60(2): 363-72.
18. Kinkead E. and Leahy H. Evaluation of the acute toxicity of selected groundwater contaminants. Harry G. Armstrong Aerospace Medical Research Lab (AAMRL-TR-87-021), 10. 1987.
19. Bogen K. and Hall L. Pharmacokinetics for regulatory risk analysis: The case of 1,1,1-trichloroethane (methyl chloroform). . *Regul Toxicol Pharmacol*. 1989; 10: 26-50.
20. Albee RR. Spencer PJ, Johnson KA, et al., Lack of trigeminal nerve toxicity in rats exposed to trichloroethylene vapor for 13 weeks. *Int J Toxicology*, 2006; 25: 531-540.
21. Kumar P, Prasad AK, Mani U, et al. , Trichloroethylene induced testicular toxicity in rats exposed by inhalation. *Hum Exp Toxicology*. 2001; 20: 585-589.
22. Baillie-Hamilton PF. Chemical toxins: a hypothesis to explain the global obesity epidemic. *Journal of alternative and complementary medicine*. 2002; 8(2): 185-92.
23. Grun F. and Blumberg B. Environmental obesogens: organotins and endocrine disruption via nuclear receptor signaling. *Endocrinology*. 2006, 147: 50-5.
24. Heindel J. Endocrine disruptors and the obesity epidemic. *Toxicol Sci*. 2003; 76(2): 247-249. .
25. Holtcamp W, Obesogens: Environmental link to obesity?. *Environ Health Perspect*. 2012; 120(2): 62-68.
26. Newbold RR., et al., Developmental exposure to endocrine disruptors and the obesity epidemic. *Reproductive toxicology*. 2007; 23(3): 290-6.

Fertility and Reproductive Outcome in Mice Following ...

27. Pelletier C, Imbeault P, Tremblay A. Energy balance and pollution by organochlorines and polychlorinated biphenyls. *Obes Rev.* 2003; 4 (1): 17-24.
28. Pise V. and Reigle T. Effects of acute inhalation exposure to 1,1,1-trichloroethane on the hypothalamo-pituitary-adrenal axis in male Sprague-Dawley rats. *J Toxicol Environ Health.* 1998; 193-208.
29. Adams, E, Spencer H, and Rowe V, et al., Vapor toxicity of 1,1,1-trichloroethane (methylchloroform) determined by experiments on laboratory animals. . *Arch Ind Hyg Occup Med.* 1950; (2): 225-236.
30. Hutcheon D, Dapson S, Gilani S, Hutcheon DE, Dapson S, Gilani SH. Persistent ductus arteriosus in weanling rats maternally exposed to methyl chloroform. *Vasc Surg.* 1985; 19:299.
31. Dapson S, Hutcheon D, and Lehr D. Effect of methyl chloroform on cardiovascular development in rats. *Teratology.* 1984; 29: 25.
32. Calhoun L, Quast F, and Schumann A et al. , Chloroethene VG: Preliminary studies to establish exposure concentrations for a chronic inhalation study with rats and mice. Midland, MI: Health and Environmental Sciences, The Dow Chemical Company. 1981.
33. Eben A, and Kimmerle G. Metabolism, excretion and toxicology of methylchloroform in acute and subacute exposed rats . *Arch Toxicol.* 1974; 31: 233-242.
34. Torkelson T, Oyen F, and McCollister D. Toxicity of 1,1,1-trichloroethane as determined on laboratory animals and human subjects. . *Am Ind Hyg Assoc J.* 1958; 19: 353-362.
35. Truffert L, Girard-Wallon C, and Emmerich E et al. , Early experimental demonstration of the hepatotoxicity of some chlorinated solvents by the study of the synthesis of hepatic DNA. *Arch Mal Prof.* 1977; 38: 261-263.
36. Lane R, Riddle B, and Borzelleca J. Effects of 1,2-dichloroethane and 1,1,1-trichloroethane in drinking water on reproduction and development in mice. *Toxicol Appl Pharmacol.* 1982; 63:409-421.
37. NTP. Final report part 1. Developmental toxicity evaluation of 1,1,1-trichloroethane (CAS No. 71-55-6) administered to CD rats. Research Triangle Park, NC: National Toxicology Program. PB88131321, 1988a.
38. George J, Price C, and Marr MC. Developmental toxicity of 1,1,1-trichloroethane in CD rats. *Fundam Appl Toxicol.* 1989;13: 641-651.
39. Van-Larebeke NA, Sasco AJ, Brophy JT, Keith MM, Gilbertson M, Watterson A. Sex ratio changes as sentinel health events of endocrine disruption. *International Journal of Occupational and Environmental Health.* 2008; 14: 138-143.
40. Hertz-Picciotto I, Jusko T, Willman EJ, Baker R, Keller JA, Teplin SW, Charles MJ. A cohort study of in utero polychlorinated biphenyl (PCB) exposures in relation to secondary sex ratio. *Environmental Health,* 2008; 7.
41. Ishihara K, Ohsako S, Tasaka K, Harayama H, Miyake M, Warita K, Tanida T, Mitsuhashi T, Nanmori T, Tabuchi Y, Yokoyama T, Kitagawa H, Hoshi N. When does the sex ratio of offspring of the paternal 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) exposure decrease: In the spermatozoa stage or at fertilization?. *Reproductive Toxicology.* 2010; 29: 68-73.